

## WEST Search History

DATE: Monday, June 24, 2002

Set Name Query  
side by side

Hit Count Set Name  
result set

*DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=OR*

L7	L6 and @ad<19981226	11	L7
L6	(HLA adj E) and (nk or CD94\$9)	17	L6
L5	L3 and (HLA adj E)	3	L5
L4	L3 and HLAE	0	L4
L3	L2 and (nk or CD94\$9)	70	L3
L2	(Braud)[IN] OR (Allan) or (ogg)[in] or (ocallaghan)[in] or (mcmichael)[in]	32981	L2
L1	(Braud)[IN] OR (Allan)	32236	L1

END OF SEARCH HISTORY

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal644axd

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\*\*\*\*\* Welcome to STN International \*\*\*\*\*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Jan 25 BLAST(R) searching in REGISTRY available in STN on the Web  
NEWS 3 Jan 29 FSTA has been reloaded and moves to weekly updates  
NEWS 4 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update frequency  
NEWS 5 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02  
NEWS 6 Mar 08 Gene Names now available in BIOSIS  
NEWS 7 Mar 22 TOXLIT no longer available  
NEWS 8 Mar 22 TRCTHERMO no longer available  
NEWS 9 Mar 28 US Provisional Priorities searched with P in CA/Caplus and USPATFULL  
NEWS 10 Mar 28 LIPINSKI/CALC added for property searching in REGISTRY  
NEWS 11 Apr 02 PAPERCHEM no longer available on STN. Use PAPERCHEM2 instead.  
NEWS 12 Apr 08 "Ask CAS" for self-help around the clock  
NEWS 13 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area  
NEWS 14 Apr 09 ZDB will be removed from STN  
NEWS 15 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB  
NEWS 16 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS  
NEWS 17 Apr 22 BIOSIS Gene Names now available in TOXCENTER  
NEWS 18 Apr 22 Federal Research in Progress (FEDRIP) now available  
NEWS 19 Jun 03 New e-mail delivery for search results now available  
NEWS 20 Jun 10 MEDLINE Reload  
NEWS 21 Jun 10 PCTFULL has been reloaded

NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,  
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),  
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*

FILE 'HOME' ENTERED AT 13:53:22 ON 24 JUN 2002

=> file medline caplus embase biosis

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	0.21	0.21

FILE 'MEDLINE' ENTERED AT 13:53:34 ON 24 JUN 2002

FILE 'CAPLUS' ENTERED AT 13:53:34 ON 24 JUN 2002

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'EMBASE' ENTERED AT 13:53:34 ON 24 JUN 2002

COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 13:53:34 ON 24 JUN 2002

COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

=> s Braud V7/au or Allan D7/au or Ogg G7/au or OCallaghan C7/au or Mcmichael A7/au  
L1 3511 BRAUD V7/AU OR ALLAN D7/AU OR OGG G7/AU OR OCALLAGHAN C7/AU OR MCMICHAEL A7/AU

=> s l3 and (nk?)

L3 NOT FOUND

The L-number entered could not be found. To see the definition of L-numbers, enter DISPLAY HISTORY at an arrow prompt (=>).

=> s l2 and (nk?)

L2 NOT FOUND

The L-number entered could not be found. To see the definition of L-numbers, enter DISPLAY HISTORY at an arrow prompt (=>).

=> s l1 and (nk?)

L2 47 L1 AND (NK?)

=> s l2 and (HLA (1N) E)

L3 35 L2 AND (HLA (1N) E)

=> dup rem l3

PROCESSING COMPLETED FOR L3

L4 14 DUP REM L3 (21 DUPLICATES REMOVED)

=> dis l4 1-14 ibib abs kwic

L4	ANSWER 1 OF 14	MEDLINE	DUPLICATE 1
ACCESSION NUMBER:	2002292488	IN-PROCESS	
DOCUMENT NUMBER:	22028985	PubMed ID: 12032324	
TITLE:	UL40-mediated NK evasion during productive		

infection with human cytomegalovirus.  
 AUTHOR: Wang Eddie C Y; McSharry Brian; Retiere Christelle; Tomasec Peter; Williams Sheila; Borysiewicz Leszek K; Braud Veronique M; Wilkinson Gavin W G  
 CORPORATE SOURCE: Section of Infection and Immunity, University of Wales College of Medicine, Tenovus Building, Heath Park, Cardiff CF14 4XX, United Kingdom.  
 SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (2002 May 28) 99 (11) 7570-5. Journal code: 7505876. ISSN: 0027-8424.  
 PUB. COUNTRY: United States  
 LANGUAGE: English  
 FILE SEGMENT: IN-PROCESS; NONINDEXED; Priority Journals  
 ENTRY DATE: Entered STN: 20020529  
 Last Updated on STN: 20020529

AB Human cytomegalovirus (HCMV) exploits a range of strategies to evade and modulate the immune response. Its capacity to down-regulate MHC I expression was anticipated to render infected cells vulnerable to natural killer (NK) attack. Kinetic analysis revealed that during productive infection, HCMV strain AD169 first enhanced and then inhibited lysis of primary skin fibroblasts by a CD94/NKG2A(+) NK cell line. The inhibition of cytotoxicity against strain AD169-infected fibroblasts was abolished by prior treatment of targets or effectors with anti-MHC I and anti-CD94 monoclonal antibodies, respectively, implying a CD94/HLA-E-dependent mechanism. An HCMV strain AD169, UL40 deletion mutant could not inhibit CD94/NKG2A(+) NK killing against skin fibroblasts. The contribution of UL40 to evasion of primary NK cells then was tested in a system where targets and effectors were MHC-matched. Primary NK cells activated with IFNalpha as well as cultured primary NK cell lines showed increased killing against DeltaUL40-infected fibroblasts compared with AD169-infected targets. This effect was abrogated by depletion of CD94(+) cells. These findings demonstrate that HCMV encodes a mechanism of evasion specifically targeted against a proportion of CD94(+) NK cells and show that this system functions during a productive infection.

TI UL40-mediated NK evasion during productive infection with human cytomegalovirus.

AU Wang Eddie C Y; McSharry Brian; Retiere Christelle; Tomasec Peter; Williams Sheila; Borysiewicz Leszek K; Braud Veronique M; Wilkinson Gavin W G

AB . . . the immune response. Its capacity to down-regulate MHC I expression was anticipated to render infected cells vulnerable to natural killer (NK) attack. Kinetic analysis revealed that during productive infection, HCMV strain AD169 first enhanced and then inhibited lysis of primary skin fibroblasts by a CD94/NKG2A(+) NK cell line. The inhibition of cytotoxicity against strain AD169-infected fibroblasts was abolished by prior treatment of targets or effectors with anti-MHC I and anti-CD94 monoclonal antibodies, respectively, implying a CD94/HLA-E-dependent mechanism. An HCMV strain AD169, UL40 deletion mutant could not inhibit CD94/NKG2A(+) NK killing against skin fibroblasts. The contribution of UL40 to evasion of primary NK cells then was tested in a system where targets and effectors were MHC-matched. Primary NK cells activated with IFNalpha as well as cultured primary NK cell lines showed increased killing against DeltaUL40-infected fibroblasts compared with AD169-infected targets. This effect was abrogated by depletion of CD94(+) cells. These findings demonstrate that HCMV encodes a mechanism of evasion specifically targeted against a proportion of CD94(+) NK cells and show that this system functions during a productive infection.

L4 ANSWER 2 OF 14 MEDLINE DUPLICATE 2  
 ACCESSION NUMBER: 2002187376 MEDLINE  
 DOCUMENT NUMBER: 21916945 PubMed ID: 11920559  
 TITLE: Human T cell receptor-mediated recognition of HLA-E.  
 AUTHOR: Garcia Pilar; Llano Manuel; de Heredia Agustin B; Willberg Christian B; Caparros Esther; Aparicio Pedro; Braud Veronique M; Lopez-Botet Miguel  
 CORPORATE SOURCE: DCEXS (Immunologia), Universitat Pompeu Fabra, Barcelona, Spain.  
 SOURCE: EUROPEAN JOURNAL OF IMMUNOLOGY, (2002 Apr) 32 (4) 936-44. Journal code: 1273201. ISSN: 0014-2980.  
 PUB. COUNTRY: Germany; Germany, Federal Republic of  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200205  
 ENTRY DATE: Entered STN: 20020403  
 Last Updated on STN: 20020522  
 Entered Medline: 20020520

AB The HLA-E class Ib molecule presents hydrophobic peptides derived from the leader sequences of other class I molecules, constituting the ligands for CD94/NKG2 lectin-like receptors. Along the course of our studies on human CD94+ T cells, we characterized an alpha beta CD8+CD94/NKG2C+ CTL clone (K14). In cytolytic assays against the murine TAP-deficient RMA-S cells transfected with human beta2 microglobulin and HLA-E (RMA-S/HLA-E), loaded with different synthetic peptides, K14 displayed a pattern of specific recognition distinct to that observed in CD94/NKG2C+ NK clones tested in parallel. RMA-S/HLA-E cells loaded with some but not all HLA class I leader sequence peptides were efficiently recognized by K14 but not by CD94/NKG2C clones, and vice versa. Remarkably, K14 also reacted with HLA-E loaded with a peptide derived from the B2LF-1 Epstein-Barr virus protein. Anti-CD94 mAb did not prevent K14 cytotoxicity against RMA-S/HLA-E cells, whereas incubation with anti-clonotypic mAb specific for the K14 TCR markedly inhibited lysis. Soluble HLA-E tetramers refolded with different peptides (i.e. VMAPRTVL, VMAPRTLIL, VMAPRTLFL) specifically stained K14 cells. HLA-E tetramer binding was minimally reduced by pretreatment with anti-CD94 mAb alone, but was completely prevented in combination with anti-clonotypic mAb. Altogether, the data unequivocally imply the generation of human T cells potentially recognizing through the alpha beta TCR HLA-E molecules that bind to class I- and virus-derived peptides.

TI Human T cell receptor-mediated recognition of HLA-E.

AU Garcia Pilar; Llano Manuel; de Heredia Agustin B; Willberg Christian B; Caparros Esther; Aparicio Pedro; Braud Veronique M; Lopez-Botet Miguel

AB The HLA-E class Ib molecule presents hydrophobic peptides derived from the leader sequences of other class I molecules, constituting the ligands for CD94/NKG2 lectin-like receptors. Along the course of our studies on human CD94+ T cells, we characterized an alpha beta CD8+CD94/NKG2C+ CTL clone (K14). In cytolytic assays against the murine TAP-deficient RMA-S cells transfected with human beta2 microglobulin and HLA-E (RMA-S/HLA-E), loaded with different synthetic peptides, K14 displayed a pattern of specific recognition distinct to that observed in CD94/NKG2C+ NK clones tested in parallel. RMA-S/HLA-E cells loaded with some but not all HLA class I leader sequence peptides were efficiently recognized by K14 but not by CD94/NKG2C clones, and vice versa. Remarkably, K14 also reacted with HLA-E loaded with a peptide derived from the BZLF-1 Epstein-Barr virus protein. Anti-CD94 mAb did not prevent K14 cytotoxicity against RMA-S/HLA-E cells, whereas incubation with anti-clonotypic mAb specific for the K14 TCR markedly inhibited lysis. Soluble HLA-E tetramers refolded with different peptides (i.e. VMAPRTVLL, VMAPRTLIL, VMAPRTLFL) specifically stained K14 cells. HLA-E tetramer binding was minimally reduced by pretreatment with anti-CD94 mAb alone, but was completely prevented in combination with anti-clonotypic mAb. Altogether, the data unequivocally imply the generation of human T cells potentially recognizing through the alpha beta TCR HLA-E molecules that bind to class I- and virus-derived peptides.

CN. . . (BZLF1 protein); 0 (Biopolymers); 0 (DNA-Binding Proteins); 0 (HLA Antigens); 0 (HLA-A Antigens); 0 (HLA-B Antigens); 0 (HLA-C Antigens); 0 (HLA-E antigen); 0 (Histocompatibility Antigens Class I); 0 (Membrane Glycoproteins); 0 (Peptide Fragments); 0 (Protein Sorting Signals); 0 (Receptors, Antigen, T-Cell, . . .

L4 ANSWER 3 OF 14 MEDLINE DUPLICATE 3  
 ACCESSION NUMBER: 2001669020 MEDLINE  
 DOCUMENT NUMBER: 21571709 PubMed ID: 11714810  
 TITLE: Intramembrane proteolysis of signal peptides: an essential step in the generation of HLA-E epitopes.  
 AUTHOR: Lemberg M K; Bland F A; Weihofen A; Braud V M; Martoglio B  
 CORPORATE SOURCE: Institute of Biochemistry, Swiss Federal Institute of Technology (Eidgenossische Technische Hochschule), Zurich, Switzerland.  
 SOURCE: JOURNAL OF IMMUNOLOGY, (2001 Dec 1) 167 (11) 6441-6. Journal code: 2985117R. ISSN: 0022-1767.  
 PUB. COUNTRY: United States  
 LANGUAGE: English  
 FILE SEGMENT: Abridged Index Medicus Journals; Priority Journals  
 ENTRY MONTH: 200201  
 ENTRY DATE: Entered STN: 20011121  
 Last Updated on STN: 20020124  
 Entered Medline: 20020102

AB Signal sequences of human MHC class I molecules are a unique source of epitopes for newly synthesized nonclassical HLA-E molecules. Binding of such conserved peptides to HLA-E induces its cell surface expression and protects cells from NK cell attack. After cleavage from the pre-protein, we show that the liberated MHC class I signal peptide is further processed by signal peptide peptidase in the hydrophobic, membrane-spanning region. This cut is essential for the release of the HLA-E epitope-containing fragment from the lipid bilayer and its subsequent transport into the lumen of the endoplasmic reticulum via the TAP.

TI Intramembrane proteolysis of signal peptides: an essential step in the generation of HLA-E epitopes.

AU Lemberg M K; Bland F A; Weihofen A; Braud V M; Martoglio B

AB Signal sequences of human MHC class I molecules are a unique source of epitopes for newly synthesized nonclassical HLA-E molecules. Binding of such conserved peptides to HLA-E induces its cell surface expression and protects cells from NK cell attack. After cleavage from the pre-protein, we show that the liberated MHC class I signal peptide is further processed by signal peptide peptidase in the hydrophobic, membrane-spanning region. This cut is essential for the release of the HLA-E epitope-containing fragment from the lipid bilayer and its subsequent transport into the lumen of the endoplasmic reticulum via the TAP.

CN 0 (ATP-Binding Cassette Transporters); 0 (Epitopes); 0 (HLA Antigens); 0 (HLA-A 0301 antigen); 0 (HLA-A Antigens); 0 (HLA-E antigen); 0 (Histocompatibility Antigens Class I); 0 (Membrane Proteins); 0 (Peptide Fragments); 0 (Protein Precursors); 0 (Protein Sorting Signals); 0. . .

L4 ANSWER 4 OF 14 MEDLINE DUPLICATE 4  
 ACCESSION NUMBER: 2000386466 MEDLINE  
 DOCUMENT NUMBER: 20354863 PubMed ID: 10898498  
 TITLE: HLA-E is expressed on trophoblast and interacts with CD94/NKG2 receptors on decidual NK cells.  
 AUTHOR: King A; Allan D S; Bowen M; Powis S J; Joseph S; Verma S; Hiby S E; McMichael A J; Loke Y W; Braud V M  
 CORPORATE SOURCE: Department of Pathology, University of Cambridge.. akk27@cam.ac.uk  
 SOURCE: EUROPEAN JOURNAL OF IMMUNOLOGY, (2000 Jun) 30 (6) 1623-31. Journal code: 1273201. ISSN: 0014-2980.  
 PUB. COUNTRY: GERMANY: Germany, Federal Republic of  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 200008  
 ENTRY DATE: Entered STN: 20000818  
 Last Updated on STN: 20000818  
 Entered Medline: 20000809

AB Non-classical MHC class I molecule HLA-E is the ligand for CD94/NKG2 NK cell receptors. Surface expression of HLA-E requires binding of specific HLA class I leader sequences. The uterine mucosa in early pregnancy (decidua) is infiltrated by large numbers of NK cells, which are closely associated with placental trophoblast cells. In this study we demonstrate that trophoblast cells express HLA-E on their cell surface in addition to the previously reported expression of HLA-G and HLA-C. Furthermore, we show that the vast majority of decidual NK cells bind to HLA-E tetrameric complexes and this binding is inhibited

by mAb to CD94. Thus, recognition of fetal HLA-E by decidual NK cells may play a key role in regulation of placental. The functional consequences of decidual NK cell interaction were investigated in cytotoxicity assays using polyclonal decidual NK cells. The overall effect of CD94/NKG2 interaction with HLA-E is inhibition of cytotoxicity by decidual NK cells. However, since decidual NK cells are unable to kill trophoblast even in the presence of mAb to MHC class I molecules and NK cell receptors, HLA-E interaction with CD94/NKG2 receptors may regulate other functions besides cytolysis during implantation.

TI HLA-E is expressed on trophoblast and interacts with CD94/NKG2 receptors on decidual NK cells.  
 AU King A; Allan D S; Bowen M; Powis S J; Joseph S; Verma S; Hiby S E; McMichael A J; Loke Y W; Braud V M  
 AB Non-classical MHC class I molecule HLA-E is the ligand for CD94/NKG2 NK cell receptors. Surface expression of HLA-E requires binding of specific HLA class I leader sequences. The uterine mucosa in early pregnancy (decidua) is infiltrated by large numbers of NK cells, which are closely associated with placental trophoblast cells. In this study we demonstrate that trophoblast cells express HLA-E on their cell surface in addition to the previously reported expression of HLA-G and HLA-C. Furthermore, we show that the vast majority of decidual NK cells bind to HLA-E tetrameric complexes and this binding is inhibited by mAb to CD94. Thus, recognition of fetal HLA-E by decidual NK cells may play a key role in regulation of placental. The functional consequences of decidual NK cell interaction were investigated in cytotoxicity assays using polyclonal decidual NK cells. The overall effect of CD94/NKG2 interaction with HLA-E is inhibition of cytotoxicity by decidual NK cells. However, since decidual NK cells are unable to kill trophoblast even in the presence of mAb to MHC class I molecules and NK cell receptors, HLA-E interaction with CD94/NKG2 receptors may regulate other functions besides cytolysis during implantation.  
 CN 0 (Antigens, CD); 0 (HLA Antigens); 0 (HLA-C Antigens); 0 (HLA-E antigen); 0 (HLA-G antigen); 0 (Histocompatibility Antigens Class I); 0 (Ligands); 0 (Membrane Glycoproteins); 0 (NKG2 protein); 0 (Receptors, Immunologic); 0 (antigen CD94)

L4 ANSWER 5 OF 14 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 5  
 ACCESSION NUMBER: 2000:118749 CAPLUS  
 DOCUMENT NUMBER: 132:292398  
 TITLE: Surface expression of HLA-E, an inhibitor of natural killer cells, enhanced by human cytomegalovirus gpUL40  
 AUTHOR(S): Tomasec, Peter; Braud, Veronique M.; Rickards, Carole; Powell, Martin B.; McSharry, Brian P.; Gadola, Stephan; Cerundolo, Vincenzo; Borysiewicz, Leszek K.; McMichael, Andrew J.; Wilkinson, Gavin W. G.  
 CORPORATE SOURCE: Department of Medicine, University of Wales College of Medicine, Cardiff, CF14 4XN, UK  
 SOURCE: Science (Washington, D. C.) (2000), 287(5455), 1031-1033  
 CODEN: SCIEAS; ISSN: 0036-8075  
 PUBLISHER: American Association for the Advancement of Science  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The nonclassical major histocompatibility complex (MHC) class I mol. HLA-E inhibits natural killer (NK) cell-mediated lysis by interacting with CD94/NKG2A receptors. Surface expression of HLA-E depends on binding of conserved peptides derived from MHC class I mols. The same peptide is present in the leader sequence of the human cytomegalovirus (HCMV) glycoprotein UL40 (gpUL40). It is shown that, independently of the transporter assocd. with antigen processing, gpUL40 can up-regulate expression of HLA-E, which protects targets from NK cell lysis. While classical MHC class I mols. are down-regulated, HLA-E is up-regulated by HCMV. Induction of HLA-E surface expression by gpUL40 may represent an escape route for HCMV.

REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI Surface expression of HLA-E, an inhibitor of natural killer cells, enhanced by human cytomegalovirus gpUL40  
 AU Tomasec, Peter; Braud, Veronique M.; Rickards, Carole; Powell, Martin B.; McSharry, Brian P.; Gadola, Stephan; Cerundolo, Vincenzo; Borysiewicz, Leszek K.; McMichael, Andrew J.; Wilkinson, Gavin W. G.  
 AB The nonclassical major histocompatibility complex (MHC) class I mol. HLA-E inhibits natural killer (NK) cell-mediated lysis by interacting with CD94/NKG2A receptors. Surface expression of HLA-E depends on binding of conserved peptides derived from MHC class I mols. The same peptide is present in the leader sequence of the human cytomegalovirus (HCMV) glycoprotein UL40 (gpUL40). It is shown that, independently of the transporter assocd. with antigen processing, gpUL40 can up-regulate expression of HLA-E, which protects targets from NK cell lysis. While classical MHC class I mols. are down-regulated, HLA-E is up-regulated by HCMV. Induction of HLA-E surface expression by gpUL40 may represent an escape route for HCMV.  
 IT Antigen receptors  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (CD94/NKG2A; human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)  
 IT CD antigens  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (CD94; human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)  
 IT Histocompatibility antigens  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (HLA-E; human cytomegalovirus gpUL40 upregulates

- surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)
- IT Glycoproteins, specific or class  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
(UL40; human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)
- IT Cytotoxicity  
Human herpesvirus 5  
(human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)
- IT Signal peptides  
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)  
(human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)
- IT Lymphocyte  
(natural killer cell; human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)
- IT Infection  
(viral; human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)
- IT 264585-39-9 264585-40-2 264585-41-3  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study)  
(amino acid sequence; human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)
- IT 205491-11-8  
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)  
(human cytomegalovirus gpUL40 upregulates surface expression of HLA-E, protects against lysis by CD94/NKG2A natural killer cells, and contains the same peptide as HLA-E derived from MHC class I mols.)

#### L4 ANSWER 6 OF 14

MEDLINE  
ACCESSION NUMBER: 2000134720 MEDLINE  
DOCUMENT NUMBER: 20134720 PubMed ID: 10669413  
TITLE: Surface expression of HLA-E, an inhibitor of natural killer cells, enhanced by human cytomegalovirus gpUL40.  
AUTHOR: Tomasec P; Braud V M; Rickards C; Powell M B; McSharry B P; Gadola S; Cerundolo V; Borysiewicz L K; McMichael A J; Wilkinson G W  
CORPORATE SOURCE: Department of Medicine, University of Wales College of Medicine, Cardiff CF14 4XN, UK.  
SOURCE: SCIENCE, (2000 Feb 11) 287 (5455) 1031.  
PUB. COUNTRY: Journal code: 0404511. ISSN: 0036-8075.  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 200002  
ENTRY DATE: Entered STN: 20000309  
Last Updated on STN: 20000309  
Entered Medline: 20000224

AB The nonclassical major histocompatibility complex (MHC) class I molecule HLA-E inhibits natural killer (NK) cell-mediated lysis by interacting with CD94/NKG2A receptors. Surface expression of HLA-E depends on binding of conserved peptides derived from MHC class I molecules. The same peptide is present in the leader sequence of the human cytomegalovirus (HCMV) glycoprotein UL40 (gpUL40). It is shown that, independently of the transporter associated with antigen processing, gpUL40 can up-regulate expression of HLA-E, which protects targets from NK cell lysis. While classical MHC class I molecules are down-regulated, HLA-E is up-regulated by HCMV. Induction of HLA-E surface expression by gpUL40 may represent an escape route for HCMV.

TI Surface expression of HLA-E, an inhibitor of natural killer cells, enhanced by human cytomegalovirus gpUL40.

AU Tomasec P; Braud V M; Rickards C; Powell M B; McSharry B P; Gadola S; Cerundolo V; Borysiewicz L K; McMichael A J; Wilkinson G W

AB The nonclassical major histocompatibility complex (MHC) class I molecule HLA-E inhibits natural killer (NK) cell-mediated lysis by interacting with CD94/NKG2A receptors. Surface expression of HLA-E depends on binding of conserved peptides derived from MHC class I molecules. The same peptide is present in the leader sequence of the human cytomegalovirus (HCMV) glycoprotein UL40 (gpUL40). It is shown that, independently of the transporter associated with antigen processing, gpUL40 can up-regulate expression of HLA-E, which protects targets from NK cell lysis. While classical MHC class I molecules are down-regulated, HLA-E is up-regulated by HCMV. Induction of HLA-E surface expression by gpUL40 may represent an escape route for HCMV.

CN 0 (HLA Antigens); 0 (HLA-E antigen); 0 (Histocompatibility Antigens Class I); 0 (Protein Sorting Signals); 0 (Receptors, Immunologic); 0 (Recombinant Fusion Proteins); 0 (UL40 glycoprotein); . . .

#### L4 ANSWER 7 OF 14 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

ACCESSION NUMBER: 2000:479385 BIOSIS  
DOCUMENT NUMBER: PREV200000479385  
TITLE: HLA-E is expressed on trophoblast and interacts with CD94/NKG2 receptors on decidual NK cells.  
AUTHOR(S): King, A. (1); Allan, D. S. J. (1); Bowen, J. M. (1); Powis, S. J. (1); Joseph, S. (1); Verma, S. (1); Hiby,

CORPORATE SOURCE: S. E. (1); McMichael, A. J. (1); Loke, Y. W. (1);  
Braud, V. M. (1)  
(1) Department of Pathology, University of Cambridge,  
Cambridge UK  
SOURCE: Placenta, (September, 2000) Vol. 21, No. 7, pp. A.39.  
print.  
Meeting Info.: 14th Rochester Trophoblast Conference  
Meeting in Association with the Society for the  
Investigation of Early Pregnancy and the 6th Meeting of the  
International Federation of Placental Associations  
Rochester, New York, USA October 04-08, 2000  
ISSN: 0143-4004.  
DOCUMENT TYPE: Conference  
LANGUAGE: English  
SUMMARY LANGUAGE: English

TI HLA-E is expressed on trophoblast and interacts with  
CD94/NKG2 receptors on decidual NK cells.  
AU King, A. (1); Allan, D. S. J. (1); Bowen, J. M. (1); Powis, S.  
J. (1); Joseph, S. (1); Verma, S. (1); Hiby, S. E. (1); McMichael, A.  
J. (1); Loke, Y. W. (1); Braud, V. M. (1)  
IT Major Concepts  
Immune System (Chemical Coordination and Homeostasis); Reproductive  
System (Reproduction)  
IT Chemicals & Biochemicals  
CD-94-NKG-2 receptor: HLA-E  
histocompatibility antigen interaction, decidual natural killer cell  
expression; HLA-E histocompatibility antigen:  
trophoblast expression

L4 ANSWER 8 OF 14 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1999:375752 CAPLUS  
DOCUMENT NUMBER: 131:18007  
TITLE: HLA-E binding  
INVENTOR(S): Braud, Veronique M.; Allan, David S.  
J.; Ogg, Graham S.; O'Callaghan,  
Christopher A.; McMichael, Andrew J.  
PATENT ASSIGNEE(S): Isis Innovation Limited, UK  
SOURCE: PCT Int. Appl., 45 pp.  
CODEN: PIXXD2

DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9928748	A2	19990610	WO 1998-GB3686	19981204
WO 9928748	A3	19991223		
W: JP, US				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
EP 1036327	A2	20000920	EP 1998-959027	19981204
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2002513541	T2	20020514	JP 2000-523554	19981204
			GB 1997-25764	A 19971204
			WO 1998-GB3686	W 19981204

PRIORITY APPLN. INFO.:

AB The invention relates to a method of causing an interaction of CD94/  
NKG2+ cells, which method comprises contacting the cells with  
recombinant HLA-E under binding conditions. The  
HLA-E property of binding to CD94/NKG2  
receptors on NK cells and a subset of CD8+ T cells is useful for  
targeting CD94/NKG2+ cells for a variety of purposes such as  
identification, isolation, killing or inactivation. In another word, the  
invention is useful for diagnostic and therapeutic purposes in a variety  
of conditions including cancer, lymphomas and leukemias, infections,  
prevention of fetus rejection, transplant rejection or graft-vs-host  
disease, immunodeficiency, and other autoimmune diseases (systemic lupus  
erythematosus, diabetes, thyroid diseases, vitiligo, rheumatoid arthritis,  
etc.). Thus, tetrameric complexes comprising biotinylated HLA-  
E, .beta.2m, signal sequence of HLA-B\*0801, and  
phycoerythrin-labeled extravidin were prepd. and tested for binding to  
NK cells and a subset of T cells. HLA-E  
-coated beads were prepd. for isolating CD94/NKG2+ cells, and  
recombinant biotinylated HLA-E coupled to  
perforin-linked extravidin was described for killing NK cells.  
Recombinant HLA-E with HLA-B8 leader  
sequence was generated for use in xenotransplantation.

TI HLA-E binding  
IN Braud, Veronique M.; Allan, David S. J.; Ogg,  
Graham S.; O'Callaghan, Christopher A.; McMichael, Andrew J.  
AB The invention relates to a method of causing an interaction of CD94/  
NKG2+ cells, which method comprises contacting the cells with  
recombinant HLA-E under binding conditions. The  
HLA-E property of binding to CD94/NKG2  
receptors on NK cells and a subset of CD8+ T cells is useful for  
targeting CD94/NKG2+ cells for a variety of purposes such as  
identification, isolation, killing or inactivation. In another word, the  
invention is useful for diagnostic and therapeutic purposes in a variety  
of conditions including cancer, lymphomas and leukemias, infections,  
prevention of fetus rejection, transplant rejection or graft-vs-host  
disease, immunodeficiency, and other autoimmune diseases (systemic lupus  
erythematosus, diabetes, thyroid diseases, vitiligo, rheumatoid arthritis,  
etc.). Thus, tetrameric complexes comprising biotinylated HLA-  
E, .beta.2m, signal sequence of HLA-B\*0801, and  
phycoerythrin-labeled extravidin were prepd. and tested for binding to  
NK cells and a subset of T cells. HLA-E  
-coated beads were prepd. for isolating CD94/NKG2+ cells, and  
recombinant biotinylated HLA-E coupled to  
perforin-linked extravidin was described for killing NK cells.  
Recombinant HLA-E with HLA-B8 leader  
sequence was generated for use in xenotransplantation.

ST HLA E CD94NKG2 NK T cell  
IT Receptors  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
(Biological study); PROC (Process)  
(CD94/NKG2; HLA-E binding for detecting,  
isolating, killing or inactivating CD94/NKG2+ NK  
cells and T cells and for diagnostic/therapeutic purposes in cancer,  
infection, transplantation or autoimmune disease)  
IT T cell (lymphocyte)  
(CD94/NKG2+; HLA-E binding for detecting,

isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT CD antigens  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (CD94; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Autoimmune disease  
 Diabetes mellitus  
 Immunodeficiency  
 Infection  
 Leukemia  
 Lymphoma  
 Neoplasm  
 Rheumatoid arthritis  
 Thyroid gland, disease  
 Transplant rejection  
 Vitiligo  
 (HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Gene, animal  
 Nucleic acids  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Avidins  
 RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Histocompatibility antigens  
 RL: BPN (Biosynthetic preparation); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)  
 (HLA-E; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Histocompatibility antigens  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (MHC (major histocompatibility complex), class I, receptors, NKG2; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Mammal (Mammalia)  
 (cells; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Chemistry  
 (chem. compds., testing; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Toxins  
 RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (delivery; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Avidins  
 RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (extr.; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Pregnancy  
 (fetus rejection prevention; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Embryo, animal  
 (fetus, rejection prevention; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Transplant and Transplantation  
 (graft-vs.-host reaction; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Lymphocyte  
 (natural killer cell, CD94/NKG2+; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Lupus erythematosus  
 (systemic; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT Animal tissue  
 Organ, animal  
 (xenogeneic; HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)



IT 9013-20-1, Streptavidin  
 RL: BSU (Biological study, unclassified); BUU (Biological use, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)  
 (HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

IT 58-85-5, Biotin  
 RL: BUU (Biological use, unclassified); BIOL (Biological study); USES (Uses)  
 (HLA-E binding for detecting, isolating, killing or inactivating CD94/NKG2+ NK cells and T cells and for diagnostic/therapeutic purposes in cancer, infection, transplantation or autoimmune disease)

L4 ANSWER 9 OF 14 CAPLUS COPYRIGHT 2002 ACS

ACCESSION NUMBER: 1999:100537 CAPLUS  
 DOCUMENT NUMBER: 130:266111  
 TITLE: MHC class I triggering by a novel cell surface ligand costimulates proliferation of activated human T cells  
 AUTHOR(S): Agrawal, Samir; Marquet, Jeanine; Freeman, Gordon J.; Tawab, Abdul; Le Bouteiller, Philippe; Roth, Patricia; Bolton, Wade; Ogg, Graham; Bousmell, Laurence; Bensussan, Armand  
 CORPORATE SOURCE: Institut National de la Sante et de la Recherche Medicale U448, Faculte de Medecine, Henri Mondor Hospital, Creteil, Fr.  
 SOURCE: Journal of Immunology (1999), 162(3), 1223-1226  
 CODEN: JOIMA3; ISSN: 0022-1767  
 PUBLISHER: American Association of Immunologists  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB BY55 is a human cell surface mol. whose expression is restricted to NK cells, a subset of circulating CD8+ T lymphocytes, and all intestinal intraepithelial T lymphocytes. Here, the authors report that BY55 is a novel NK receptor showing broad specificity for both classical and nonclassical MHC class I mols., and that optimal binding requires a prior aggregation of MHC class I complexes. Using BY55 transfectants, the authors have identified functional consequences of MHC class I/ligand interactions for the class I-bearing cell. The triggering of MHC class I mols. on human T cell clones by BY55 delivered a potent proliferative signal in the presence of sol. CD3 mAb. The costimulatory signal provided by MHC class I ligation was only seen in activated, and not resting, peripheral blood T cells. This observation represents an addnl. and/or alternative pathway to CD28 costimulation and may be of particular relevance in memory T cells lacking CD28, such as intestinal intraepithelial T lymphocytes, which are CD28- but BY55+.

REFERENCE COUNT: 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

AU Agrawal, Samir; Marquet, Jeanine; Freeman, Gordon J.; Tawab, Abdul; Le Bouteiller, Philippe; Roth, Patricia; Bolton, Wade; Ogg, Graham; Bousmell, Laurence; Bensussan, Armand

AB BY55 is a human cell surface mol. whose expression is restricted to NK cells, a subset of circulating CD8+ T lymphocytes, and all intestinal intraepithelial T lymphocytes. Here, the authors report that BY55 is a novel NK receptor showing broad specificity for both classical and nonclassical MHC class I mols., and that optimal binding requires a prior aggregation of MHC class I complexes. Using BY55 transfectants, the authors have identified functional consequences of MHC class I/ligand interactions for the class I-bearing cell. The triggering of MHC class I mols. on human T cell clones by BY55 delivered a potent proliferative signal in the presence of sol. CD3 mAb. The costimulatory signal provided by MHC class I ligation was only seen in activated, and not resting, peripheral blood T cells. This observation represents an addnl. and/or alternative pathway to CD28 costimulation and may be of particular relevance in memory T cells lacking CD28, such as intestinal intraepithelial T lymphocytes, which are CD28- but BY55+.

IT Histocompatibility antigens  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
 (HLA-E; BY55 as ligand for)

L4 ANSWER 10 OF 14 MEDLINE MEDLINE DUPLICATE 6

ACCESSION NUMBER: 1999158668 MEDLINE  
 DOCUMENT NUMBER: 99158668 PubMed ID: 10047540  
 TITLE: Functions of nonclassical MHC and non-MHC-encoded class I molecules.  
 AUTHOR: Braud V M; Allan D S; McMichael A J  
 CORPORATE SOURCE: Institute of Molecular Medicine, John Radcliffe Hospital, Oxford, UK. vbraud@worf.molbiol.ox.ac.uk  
 SOURCE: CURRENT OPINION IN IMMUNOLOGY, (1999 Feb) 11 (1) 100-8.  
 Ref: 85  
 Journal code: 8900118. ISSN: 0952-7915.  
 PUB. COUNTRY: ENGLAND: United Kingdom  
 Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 199903  
 ENTRY DATE: Entered STN: 19990326  
 Last Updated on STN: 19990326  
 Entered Medline: 19990317

AB Fascinating recent discoveries have focused attention on the nonclassical class I molecules. They can exert their function at most levels of the immune response, being part of both innate and adaptive immune systems. They not only have specialized antigen-presentation functions but also play important immunoregulatory roles. HLA-E regulates natural killer cells by interacting with CD94/NKG2 receptors; the MIC (MHC class I chain related) glycoproteins appear crucial to the activation of gamma delta T cells in the gastrointestinal epithelium; HLA-G may play a role in controlling the immune response to the fetus; and CD1 molecules are important in defense against bacterial infections, as well as in the development and regulation of a subset of NKT cells expressing a highly restricted TCR repertoire; however not all nonclassical class I molecules have an immunological function, as demonstrated by HFE which is implicated in iron metabolism.

AU Braud V M; Allan D S; McMichael A J  
 AB . . . of both innate and adaptive immune systems. They not only have specialized antigen-presentation functions but also play important immunoregulatory roles. HLA-E regulates natural killer

cells by interacting with CD94/NKG2 receptors; the MIC (MHC class I chain related) glycoproteins appear crucial to the activation of gamma delta T cells in the . . . molecules are important in defense against bacterial infections, as well as in the development and regulation of a subset of NKT cells expressing a highly restricted TCR repertoire; however not all nonclassical class I molecules have an immunological function, as demonstrated. . .

CN 0 (Antigens, CD); 0 (HLA Antigens); 0 (HLA-E antigen); 0 (HLA-G antigen); 0 (HLA-H antigen); 0 (Histocompatibility Antigens Class I); 0 (MICA protein); 0 (MICB antigen); 0 (Q. . .

L4 ANSWER 11 OF 14 MEDLINE DUPLICATE 7  
 ACCESSION NUMBER: 1999383027 MEDLINE  
 DOCUMENT NUMBER: 99383027 PubMed ID: 10453651  
 TITLE: Regulation of NK cell functions through interaction of the CD94/NKG2 receptors with the nonclassical class I molecule HLA-E.  
 AUTHOR: Braud V M; McMichael A J  
 CORPORATE SOURCE: Institute of Molecular Medicine, Headington, Oxford, UK.  
 SOURCE: CURRENT TOPICS IN MICROBIOLOGY AND IMMUNOLOGY, (1999) 244 85-95. Ref: 30  
 Journal code: 0110513. ISSN: 0070-217X.  
 PUB. COUNTRY: GERMANY: Germany, Federal Republic of  
 Journal; Article; (JOURNAL ARTICLE)  
 General Review; (REVIEW)  
 (REVIEW, TUTORIAL)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 199910  
 ENTRY DATE: Entered STN: 19991014  
 Last Updated on STN: 19991014  
 Entered Medline: 19991006

TI Regulation of NK cell functions through interaction of the CD94/NKG2 receptors with the nonclassical class I molecule HLA-E.

AU Braud V M; McMichael A J  
 CN 0 (Antigens, CD); 0 (HLA Antigens); 0 (HLA-E antigen); 0 (Histocompatibility Antigens Class I); 0 (Ligands); 0 (Membrane Glycoproteins); 0 (NKG2 protein); 0 (Peptides); 0 (Receptors, Immunologic); 0 (antigen CD94)

L4 ANSWER 12 OF 14 MEDLINE DUPLICATE 8  
 ACCESSION NUMBER: 1998146055 MEDLINE  
 DOCUMENT NUMBER: 98146055 PubMed ID: 9486650  
 TITLE: HLA-E binds to natural killer cell receptors CD94/NKG2A, B and C.  
 COMMENT: Comment in: Nature. 1998 Feb 19;391(6669):740-1, 743  
 AUTHOR: Braud V M; Allan D S; O'Callaghan C A; Soderstrom K; D'Andrea A; Ogg G S; Lazetic S; Young N T; Bell J I; Phillips J H; Lanier L L; McMichael A J  
 CORPORATE SOURCE: Institute of Molecular Medicine, John Radcliffe Hospital, Oxford, UK. vbraud@wolf.molbiol.ox.ac.uk  
 SOURCE: NATURE, (1998 Feb 19) 391 (6669) 795-9.  
 Journal code: 0410462. ISSN: 0028-0836.  
 PUB. COUNTRY: ENGLAND: United Kingdom  
 Journal; Article; (JOURNAL ARTICLE)  
 LANGUAGE: English  
 FILE SEGMENT: Priority Journals  
 ENTRY MONTH: 199803  
 ENTRY DATE: Entered STN: 19980319  
 Last Updated on STN: 19980319  
 Entered Medline: 19980309

AB The protein HLA-E is a non-classical major histocompatibility complex (MHC) molecule of limited sequence variability. Its expression on the cell surface is regulated by the binding of peptides derived from the signal sequence of some other MHC class I molecules. Here we report the identification of ligands for HLA-E. We constructed tetramers in which recombinant HLA-E and beta2-microglobulin were refolded with an MHC leader-sequence peptide, biotinylated, and conjugated to phycoerythrin-labelled Extravidin. This HLA-E tetramer bound to natural killer (NK) cells and a small subset of T cells from peripheral blood. On transfectants, the tetramer bound to the CD94/NKG2A, CD94/NKG2B and CD94/NKG2C NK cell receptors, but did not bind to the immunoglobulin family of NK cell receptors (KIR). Surface expression of HLA-E was enough to protect target cells from lysis by CD94/NKG2A+ NK-cell clones. A subset of HLA class I alleles has been shown to inhibit killing by CD94/NKG2A+ NK-cell clones. Only the HLA alleles that possess a leader peptide capable of upregulating HLA-E surface expression confer resistance to NK-cell-mediated lysis, implying that their action is mediated by HLA-E, the predominant ligand for the NK cell inhibitory receptor CD94/NKG2A.

TI HLA-E binds to natural killer cell receptors CD94/NKG2A, B and C.

AU Braud V M; Allan D S; O'Callaghan C A; Soderstrom K; D'Andrea A; Ogg G S; Lazetic S; Young N T; Bell J I; Phillips J H; Lanier L L; McMichael A J

AB The protein HLA-E is a non-classical major histocompatibility complex (MHC) molecule of limited sequence variability. Its expression on the cell surface is regulated. . . derived from the signal sequence of some other MHC class I molecules. Here we report the identification of ligands for HLA-E. We constructed tetramers in which recombinant HLA-E and beta2-microglobulin were refolded with an MHC leader-sequence peptide, biotinylated, and conjugated to phycoerythrin-labelled Extravidin. This HLA-E tetramer bound to natural killer (NK) cells and a small subset of T cells from peripheral blood. On transfectants, the tetramer bound to the CD94/NKG2A, CD94/NKG2B and CD94/NKG2C NK cell receptors, but did not bind to the immunoglobulin family of NK cell receptors (KIR). Surface expression of HLA-E was enough to protect target cells from lysis by CD94/NKG2A+ NK-cell clones. A subset of HLA class I alleles has been shown to inhibit killing by CD94/NKG2A+ NK-cell clones. Only the HLA alleles that possess a leader peptide capable of upregulating HLA-E surface expression confer resistance to NK-cell-mediated lysis, implying that their action is mediated by HLA-E, the predominant ligand for the NK cell inhibitory receptor CD94/NKG2A.

CN 0 (Antigens, CD); 0 (HLA Antigens); 0 (HLA-E antigen);  
0 (Histocompatibility Antigens Class I); 0 (Ligands); 0 (Membrane  
Glycoproteins); 0 (NKG2 protein); 0 (Protein Sorting Signals); 0  
(Receptors, Immunologic); 0 (Recombinant Proteins); 0 (antigen CD94); 0  
(beta 2-Microglobulin); 0 (killer inhibitory).

L4 ANSWER 13 OF 14 MEDLINE DUPLICATE 9  
ACCESSION NUMBER: 1998325367 MEDLINE  
DOCUMENT NUMBER: 98325367 PubMed ID: 9660937  
TITLE: Structural features impose tight peptide binding  
specificity in the nonclassical MHC molecule HLA-  
E.  
AUTHOR: O'Callaghan C A; Tormo J; Willcox B E; Braud V M;  
Jakobsen B K; Stuart D I; McMichael A J; Bell J  
I; Jones E Y  
CORPORATE SOURCE: Nuffield Department of Clinical Medicine, University of  
Oxford, John Radcliffe Hospital, United Kingdom.  
SOURCE: MOLECULAR CELL, (1998 Mar) 1 (4) 531-41.  
Journal code: 9802571. ISSN: 1097-2765.  
PUB. COUNTRY: United States  
LANGUAGE: English  
FILE SEGMENT: Priority Journals  
ENTRY MONTH: 199807  
ENTRY DATE: Entered STN: 19980811  
Last Updated on STN: 19980811  
Entered Medline: 19980728

AB The crystal structure of the nonclassical human class Ib MHC molecule  
HLA-E has been determined in complex with a prototypic  
ligand, the nonamer peptide (VMAPRTVLL), derived from the highly conserved  
residues 3-11 of the human MHC class Ia leader sequence. The mode of  
peptide binding retains some of the standard features observed in MHC  
class Ia complexes, but novel features imply that HLA-E  
has evolved to mediate specific binding to a tightly defined set of almost  
identical hydrophobic peptides from the highly conserved class I leader  
sequences. These molecular adaptations make HLA-E a  
rigorous checkpoint at the cell surface reporting on the integrity of the  
antigen processing pathway to CD94/NKG2 receptor-bearing natural  
killer cells.

TI Structural features impose tight peptide binding specificity in the  
nonclassical MHC molecule HLA-E.

AU O'Callaghan C A; Tormo J; Willcox B E; Braud V M; Jakobsen B K;  
Stuart D I; McMichael A J; Bell J I; Jones E Y

AB The crystal structure of the nonclassical human class Ib MHC molecule  
HLA-E has been determined in complex with a prototypic  
ligand, the nonamer peptide (VMAPRTVLL), derived from the highly conserved  
residues 3-11. . . . of peptide binding retains some of the standard  
features observed in MHC class Ia complexes, but novel features imply that  
HLA-E has evolved to mediate specific binding to a  
tightly defined set of almost identical hydrophobic peptides from the  
highly conserved class I leader sequences. These molecular adaptations  
make HLA-E a rigorous checkpoint at the cell surface  
reporting on the integrity of the antigen processing pathway to CD94/  
NKG2 receptor-bearing natural killer cells.

CN 0 (HLA Antigens); 0 (HLA-B8 Antigen); 0 (HLA-E  
antigen); 0 (Histocompatibility Antigens Class I); 0 (Peptide Fragments)

L4 ANSWER 14 OF 14 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
ACCESSION NUMBER: 1999:219821 BIOSIS  
DOCUMENT NUMBER: PREV199900219821  
TITLE: Decidual NK cells have receptors for HLA  
-E which is expressed by human trophoblast.  
AUTHOR(S): Allan, D.S.J. (1); Verma, S.; Bowen, J. M.; Loke,  
Y. W.; McMichael, J. (1); Braud, V. M. (1); King,  
A.  
CORPORATE SOURCE: (1) Institute of Molecular Medicine, John Radcliffe  
Hospital, Oxford, OX3 9DS UK  
SOURCE: Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 79.  
Meeting Info.: Fifth Annual Meeting of the Society for  
Natural Immunity Seventeenth International Natural Killer  
Cell Workshop Warrenton, Virginia, USA October 17-21, 1998  
ISSN: 1018-8916.  
DOCUMENT TYPE: Conference  
LANGUAGE: English

TI Decidual NK cells have receptors for HLA-E  
which is expressed by human trophoblast.

AU Allan, D.S.J. (1); Verma, S.; Bowen, J. M.; Loke, Y. W.;  
McMichael, J. (1); Braud, V. M. (1); King, A.

IT . . .  
(Homeostasis)

IT Parts, Structures, & Systems of Organisms  
natural killer cells: blood and lymphatics, immune system

IT Chemicals & Biochemicals  
HLA-E receptor; HLA-E:  
expression

=> dui his

DUIS IS NOT A RECOGNIZED COMMAND

The previous command name entered was not recognized by the system.  
For a list of commands available to you in the current file, enter  
"HELP COMMANDS" at an arrow prompt (=>).

=> dis his

(FILE 'HOME' ENTERED AT 13:53:22 ON 24 JUN 2002)

FILE 'MEDLINE, CAPLUS, EMBASE, BIOSIS' ENTERED AT 13:53:34 ON 24 JUN 2002

L1 3511 S BRAUD V?/AU OR ALLAN D?/AU OR OGG G?/AU OR OCALLAGHAN C?/AU O  
L2 47 S L1 AND (NK?)  
L3 35 S L2 AND (HLA (1N) E)  
L4 14 DUP REM L3 (21 DUPLICATES REMOVED)

=> s nk (P) (NK? or CD94?)

L5 72958 NK (P) (NK? OR CD94?)

=> s (HLA (1N) E) (P) (NK? or CD94?)

L6 315 (HLA (1N) E) (P) (NK? OR CD94?)

=> s 16 and PD<19981204

'19981204' NOT A VALID FIELD CODE  
3 FILES SEARCHED...

=&gt; dup rem l7

PROCESSING COMPLETED FOR L7

L8 30 DUP REM L7 (33 DUPLICATES REMOVED)

=&gt; dis l8 1-30 kwic

L8 ANSWER 1 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 1

TI HLA-E is a major ligand for the natural killer

inhibitory receptor CD94/NKG2A

SO Proceedings of the National Academy of Sciences of the United States of America (1998), 95(9), 5199-5204  
CODEN: PNASA6; ISSN: 0027-8424

AB . . . a nonamer peptide derived from certain HLA class I signal sequences is a necessary requirement for the stabilization of endogenous HLA-E expression on the surface of 721.221 cells. This led the authors to examine the ability of HLA-E to protect HLA class I transfectants from natural killer (NK) cell-mediated lysis. It was possible to implicate the CD94/NKG2A complex as an inhibitory receptor recognizing this class Ib mol. by using as target a .221 transfectant selectively expressing surface HLA-E. HLA-E had no apparent inhibitory effect mediated through the identified Ig superfamily (Ig-SF) human killer cell inhibitory receptors or ILT2/LIR1. Further studies of CD94/NKG2+ NK cell-mediated recognition of .221 cells transfected with different HLA class I allotypes (i.e., -Cw4, -Cw3, -B7) confirmed that the inhibitory interaction was mediated by CD94/NKG2A recognizing the surface HLA-E mol., because only antibodies directed against either HLA-E, CD94, or CD94/NKG2A specifically restored lysis. Surface stabilization of HLA-E in cold-treated .221 cells loaded with appropriate peptides was sufficient to confer protection, resulting from recognition of the HLA class Ib mol. by the CD94/NKG2A inhibitory receptor. Consistent with the prediction that the ligand for CD94/NKG2A is expressed ubiquitously, the authors' examn. of HLA-E antigen distribution indicated that it is detectable on the surface of a wide variety of cell types.

ST HLA E ligand natural killer receptor; CD94

NKG2A receptor HLA E ligand

IT CD antigens

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(CD94, complexes with NKG2A; HLA-

E is ligand for natural killer inhibitory receptor CD94

/NKG2A)

IT Ligands

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(HLA-E is ligand for natural killer inhibitory

receptor CD94/NKG2A)

IT Histocompatibility antigens

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(HLA-E; HLA-E is ligand for

natural killer inhibitory receptor CD94/NKG2A)

IT Proteins, specific or class

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(NKG2A, complexes with CD94; HLA-

E is ligand for natural killer inhibitory receptor CD94

/NKG2A)

IT Cytolysis

(natural killer cell-mediated; HLA-E is ligand for

natural killer inhibitory receptor CD94/NKG2A)

IT Lymphocyte

(natural killer cell; HLA-E is ligand for natural

killer inhibitory receptor CD94/NKG2A)

L8 ANSWER 2 OF 30 CAPLUS COPYRIGHT 2002 ACS

SO Proceedings of the National Academy of Sciences of the United States of America (1998), 95(9), 4791-4794

CODEN: PNASA6; ISSN: 0027-8424

AB A review and discussion with 44 refs. There are abundant data to substantiate the conclusion that CD94/NKG2A receptors of human natural killer cells directly recognize HLA-E mols.

IT Histocompatibility antigens

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(HLA-E; HLA class I specificity for

natural killer cell receptor CD94/NKG2A)

L8 ANSWER 3 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 2

SO European Journal of Immunology (1998), 28(12), 4356-4361

CODEN: EJIMAF; ISSN: 0014-2980

AB Recent studies on human NK cells have demonstrated that the NK cell CD94/NKG2 receptors bind to the nonclassical MHC class I mol. HLA-E. A functional CD94/NKG2 complex has not yet been identified in rodents, but cDNA encoding rat and mouse CD94 and NKG2 have recently been cloned, suggesting that CD94/NKG2 receptors may exist in species other than man. The mouse nonclassical MHC class I mol. Qa-1 shares several features with HLA-E. This suggests that Qa-1 may be similarly recognized by murine NK cells. To study the ability of Qa-1 to bind to murine NK cells, the authors have produced a sol. tetrameric form of Qa-1b. The authors demonstrate that Qa-1b tetramers distinctly bind to a large subset of fresh or IL-2-activated NK1.1+/CD3- splenocytes independently of the expression of Ly49 inhibitory receptors. Binding occurs whether NK cells have evolved in an MHC class I-expressing or in an MHC class I-deficient environment. The data suggest the existence of a Qa-1-recognizing structure on a large subpopulation of murine NK cells that may be similar to the human CD94/NKG2 heterodimeric complex.

L8 ANSWER 4 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 3

TI HLA-E-bound peptides influence recognition by

inhibitory and triggering CD94/NKG2 receptors.

Preferential response to an HLA-G-derived nonamer

SO European Journal of Immunology (1998), 28(9), 2854-2863

- AB The HLA-E class Ib mol. constitutes a major ligand for the lectin-like CD94/NKG2 natural killer (NK) cell receptors. Specific HLA class I leader sequence-derived nonapeptides bind to endogenous HLA-E mols. in the HLA-defective cell line 721.221, inducing HLA-E surface expression, and promote CD94/NKG2-mediated recognition. The authors compared the ability of NK clones which expressed either inhibitory or activating CD94/NKG2 receptors to recognize HLA-E mols. on the surface of 721.221 cells loaded with a panel of synthetic nonamers derived from the leader sequences of most HLA class I mols. The results support the notion that the primary structure of the HLA-E-bound peptides influences CD94/NKG2-mediated recognition, beyond their ability to stabilize surface HLA-E. CD94/NKG2A+ NK clones appeared more sensitive to the interaction with most HLA-E-peptide complexes than did effector cells expressing the activating CD94/NKG2C receptor. An exception to this pattern was I-ILA-E loaded with the HLA-G-derived nonamer. This complex triggered cytotoxicity very efficiently over a wide range of peptide concns., suggesting that the HLA-E/G-nonamer complex interacts with the CD94/NKG2 triggering receptor with a higher affinity. These results raise the possibility that CD94/NKG2-mediated recognition of HLA-E expressed on extravillous cytotrophoblasts plays an important role in maternal-fetal cellular interactions.
- IT CD antigens  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(CD94; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT Receptors  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(HLA-E antigen; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT Histocompatibility antigens  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(HLA-E, Ib, complex with peptides; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT Histocompatibility antigens  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(HLA-E, receptors; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT Gene, animal  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(HLA-G; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT Receptors  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(NKG2; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT Peptides, biological studies  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(complex with HLA-E class Ib; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT Trophoblast  
(cytotrophoblast; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors in relation to)
- IT Lymphocyte  
(natural killer cell; HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- IT 193002-77-6 202657-59-8 202657-60-1 202657-61-2 202657-62-3  
205491-11-8 214621-61-1 215439-93-3 215439-97-7 215440-00-9  
215440-04-3  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors)
- L8 ANSWER 5 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 4
- TI Specific engagement of the CD94/NKG2-A killer inhibitory receptor by the HLA-E class Ib molecule induces SHP-1 phosphatase recruitment to tyrosine-phosphorylated NKG2-A. Evidence for receptor function in heterologous transfectants
- SO European Journal of Immunology (1998), 28(4), 1280-1291
- AB CODEN: EJIMAF; ISSN: 0014-2980  
It was recently demonstrated that the CD94/NKG2-A killer inhibitory receptor (KIR) specifically recognizes the HLA-E class Ib mol. The apparent CD94-mediated specific recognition of different HLA class Ia allotypes, transfected into the HLA-defective cell line 721.221, indeed depends on their selective ability to concomitantly stabilize the surface expression of endogenous HLA-E mols., which confer protection against CD94/NKG2-A+ effector cells. The authors show that a selective engagement of the CD94/NKG2-A inhibitory receptor with a specific monoclonal antibody (mAb) (Z199) was sufficient to induce Tyr phosphorylation of the NKG2-A subunit and SHP-1 recruitment. These early biochem. events, commonly related to neg. signaling pathways, were also detected upon the specific interaction of NK cells with an HLA-E+ 721.221 transfectant (.221-AEH), and were prevented by pre-incubation of .221-AEH with an anti-HLA class I mAb. MAb crosslinking of the CD94/NKG2-A receptor, segregated from other NK-assocd. mols. by transfection into a rat basophilic leukemia cell line (RBL-2H3), promoted Tyr phosphorylation of NKG2-A and co-pptn. of SHP-1, together

with an inhibition of secretory events triggered via Fc.epsilon.RI.  
Interaction of CD94/NKG2-A+ RBL cells with the  
HLA-E\*221-AEH transfectant specifically induced a  
detectable assocn. of SHP-1 with NKG2-A, constituting a more  
formal evidence for the receptor-HLA class I interaction.

ST CD94 NKG2A phosphorylation HLA E;  
SHP1 phosphatase CD94 NKG2A receptor

IT CD antigens  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological  
process); BSU (Biological study, unclassified); BIOL (Biological study);  
PROC (Process)  
(CD94, complexes, with NKG2-A; HLA-  
E ligand-induced recruitment of SHP-1 phosphatase to  
ITIM-phosphorylated NKG2-A of killer inhibitory receptor)

IT Histocompatibility antigens  
RL: BSU (Biological study, unclassified); BIOL (Biological study)  
(HLA, class I, receptors, KIR, CD94/NKG2-A;  
HLA-E ligand-induced recruitment of SHP-1 phosphatase  
to ITIM-phosphorylated NKG2-A of killer inhibitory receptor)

IT Signal transduction, biological  
(HLA-E ligand-induced recruitment of SHP-1  
phosphatase to ITIM-phosphorylated NKG2-A of killer  
inhibitory receptor)

IT Histocompatibility antigens  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological  
study, unclassified); BIOL (Biological study)  
(HLA-E; HLA-E ligand-induced  
recruitment of SHP-1 phosphatase to ITIM-phosphorylated NKG2  
-A of killer inhibitory receptor)

IT Protein motifs  
(ITIM (immunoreceptor tyrosine-based inhibitory motif); HLA-  
E ligand-induced recruitment of SHP-1 phosphatase to  
ITIM-phosphorylated NKG2-A of killer inhibitory receptor)

IT Antigen receptors  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological  
process); BSU (Biological study, unclassified); BIOL (Biological study);  
PROC (Process)  
(KIR (killer cell inhibitory), CD94/NKG2-A;  
HLA-E ligand-induced recruitment of SHP-1 phosphatase  
to ITIM-phosphorylated NKG2-A of killer inhibitory receptor)

IT Proteins, specific or class  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological  
process); BSU (Biological study, unclassified); BIOL (Biological study);  
PROC (Process)  
(NKG2-A, complexes, with CD94; HLA-  
E ligand-induced recruitment of SHP-1 phosphatase to  
ITIM-phosphorylated NKG2-A of killer inhibitory receptor)

IT Lymphocyte  
(natural killer cell; HLA-E ligand-induced  
recruitment of SHP-1 phosphatase to ITIM-phosphorylated NKG2  
-A of killer inhibitory receptor)

IT Phosphorylation, biological  
(receptor; HLA-E ligand-induced recruitment of  
SHP-1 phosphatase to ITIM-phosphorylated NKG2-A of killer  
inhibitory receptor)

IT 79747-53-8, Tyrosine phosphatase  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
(Biological study); PROC (Process)  
(SHP-1; HLA-E ligand-induced recruitment of SHP-1  
phosphatase to ITIM-phosphorylated NKG2-A of killer  
inhibitory receptor)

L8 ANSWER 6 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 5  
SO Journal of Experimental Medicine (1998), 188(5), 973-978  
CODEN: JEMEAU; ISSN: 0022-1007

AB . . . binds with high affinity and accounts for almost all of the  
peptides assocd. with this mol. Human histocompatibility leukocyte  
antigen (HLA)-E, a homolog of Qa-1b, binds similar  
peptides derived from human class Ia mols. and interacts with CD94  
/NKG2 receptors on natural killer cells. The authors used  
surface plasmon resonance to det. the ability of Qa-1b to bind related. .  
. peptides derived from the leaders of class I mols. from several  
mammalian species. All of the peptides reported to bind HLA-  
E bound readily to Qa-1b. In addn., peptides derived from leader  
segments of different mammals also bound to Qa-1b, indicating a . . .

L8 ANSWER 7 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 6  
TI Recognition of human histocompatibility leukocyte antigen (HLA)-  
E complexed with HLA class I signal sequence-derived peptides by  
CD94/NKG2 confers protection from natural killer  
cell-mediated lysis

SO Journal of Experimental Medicine (1998), 187(5), 813-818  
CODEN: JEMEAU; ISSN: 0022-1007

AB Human histocompatibility leukocyte antigen (HLA)-E is  
a nonclassical HLA class I mol., the gene for which is transcribed in most  
tissues. It has recently been. . . reported that this mol. binds  
peptides derived from the signal sequence of HLA class I proteins;  
however, no function for HLA-E has yet been described.  
The authors show that natural killer (NK) cells can recognize  
target cells expressing HLA-E mols. on the cell  
surface and this interaction results in inhibition of the lytic process.  
Furthermore, HLA-E recognition is mediated primarily  
through the CD94/NKG2-A heterodimer, as CD94  
-specific, but not killer cell inhibitory receptor (KIR)-specific mAbs  
block HLA-E-mediated protection of target cells. Cell  
surface HLA-E could be increased by incubation with  
synthetic peptides corresponding to residues 3-11 from the signal  
sequences of a no. of. . . HLA class I mols.; however, only peptides  
which contained a Met at position 2 were capable of conferring resistance  
to NK-mediated lysis, whereas those having Thr at position 2 had  
no effect. Interestingly, HLA class I mols. previously correlated with  
CD94/NKG2 recognition all have Met at residue 4 of the  
signal sequence (position 2 of the HLA-E binding  
peptide) whereas those which have been reported not to interact with  
CD94/NKG2 have Thr at this position. These data thus  
show a function for HLA-E and suggest an alternative  
explanation for the apparent broad reactivity of CD94/  
NKG2 with HLA class I mols.; that CD94/NKG2  
interacts with HLA-E complexed with signal sequence  
peptides derived from "protective" HLA class I alleles rather than  
directly interacting with classical HLA class. . .

IT Receptors

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(CD94/NKG2; recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

IT CD antigens  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(CD94; recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

IT Histocompatibility antigens  
RL: BSU (Biological study, unclassified); BIOL (Biological study) (HLA, class I; recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

IT Histocompatibility antigens  
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)  
(HLA-E; recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

IT Proteins, specific or class  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(NKG2; recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

IT Lymphocyte  
(natural killer cell; recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

IT Cytolysis  
(recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

IT 193002-77-6 202657-62-3 202657-64-5 205491-11-8  
RL: PRP (Properties)  
(recognition of HLA-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis)

L8 ANSWER 8 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 7

TI HLA-E binds to natural killer cell receptors CD94/NKG2A, B and C

SO Nature (London) (1998), 391(6669), 795-799  
CODEN: NATUAS; ISSN: 0028-0836

AB The protein HLA-E is a non-classical major histocompatibility complex (MHC) mol. of limited sequence variability. Its expression on the cell surface is regulated. . . . derived from the signal sequence of some other MHC class I mols. Here we report the identification of ligands for HLA-E. We constructed tetramers in which recombinant HLA-E and .beta.2-microglobulin were refolded with an MHC, leader-sequence peptide, biotinylated, and conjugated to phycoerythrin-labeled Extravidin. This HLA-E tetramer bound to natural killer (NK) cells and a small subset of T cells from peripheral blood. On transfectants, the tetramer bound to the CD94/NKG2A, CD94/NKG2B and CD94/NKG2C NM cell receptors, but did not bind to the Ig family of NK cell receptors (KIR). Surface expression of HLA-E was enough to protect target cells from lysis by CD94/NKG2A + NK-cell clones. A subset of HLA class I alleles has been shown to inhibit killing by CD94/NKG2A+ NK-cell clones. Only the HLA alleles that possess a leader peptide capable of upregulating HLA-E surface expression confer resistance to NK-cell-mediated lysis, implying that their action is mediated by HLA-E, the predominant ligand for the NK cell inhibitory receptor CD94/NKG2A.

ST HLA E natural killer cell receptor; CD94 NKG2 receptor natural killer cell

IT CD antigens  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(CD94; HLA-E antigen binding to natural killer cell receptors CD94/NKG2A, B and C and resistance to NK-mediated lysis)

IT Histocompatibility antigens  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(HLA, class I, alleles; HLA-E antigen binding to natural killer cell receptors CD94/NKG2A, B and C and resistance to NK-mediated lysis by leader peptides from)

IT Cytolysis  
(HLA-E antigen binding to natural killer cell receptors CD94/NKG2A, B and C and resistance to NK-mediated lysis)

IT Histocompatibility antigens  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(HLA-E; HLA-E antigen binding to natural killer cell receptors CD94/NKG2A, B and C and resistance to NK-mediated lysis)

IT Antigen receptors  
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(NKG2, variants A and B and C; HLA-E antigen binding to natural killer cell receptors CD94/NKG2A, B and C and resistance to NK-mediated lysis)

IT Lymphocyte  
(natural killer cell; HLA-E antigen binding to natural killer cell receptors CD94/NKG2A, B and C)

and resistance to NK-mediated lysis)

- L8 ANSWER 9 OF 30 CAPLUS COPYRIGHT 2002 ACS  
SO Cell (Cambridge, Massachusetts) (1998), 92(6), 705-707  
CODEN: CELLS5; ISSN: 0092-8674
- AB . . . . . discussing the function of mouse Ly49 receptors, the structure and function of human KIR receptors, and the mol. interaction of HLA-E with CD94/NKG2 heterodimers.
- IT Histocompatibility antigens  
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)  
(HLA-E; structure and function of NK cell receptors for classical and nonclassical MHC class I)
- L8 ANSWER 10 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 8  
SO Immunity (1998), 8(6), 693-701  
CODEN: IUNIEH; ISSN: 1074-7613
- AB While the inhibitory NK cell receptors for MHC class I express immunoreceptor tyrosine-based inhibitory motifs that recruit intracellular tyrosine phosphatases and prevent NK cell effector function, the activating NK cell receptors lack intrinsic sequences required for cellular stimulation. CD94/NKG2C, an activating NK cell receptor of the C-type lectin superfamily that binds to HLA-E, noncovalently assoc. with DAP12, a membrane receptor contg. an immunoreceptor tyrosine-based activating motif. Efficient expression of CD94/NKG2C on the cell surface requires the presence of DAP12, and charged residues in the transmembrane domains of DAP12 and NKG2C are necessary for this interaction. These results provide a mol. basis for the assembly of NK cell receptors for MHC class I involved in cellular activation and inhibition.
- IT Histocompatibility antigens  
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(HLA-E, receptors, CD94/NKG2C; DAP12 protein assocn. with activating CD94/NKG2C NK cell receptors)
- L8 ANSWER 11 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 9  
SO Molecular Cell (1998), 1(4), 531-541  
CODEN: MOCEFL; ISSN: 1097-2765
- AB The crystal structure of the nonclassical human class Ib MHC mol. HLA-E has been detd. in complex with a prototypic ligand, the nonamer peptide (VMAPRTVLL), derived from the highly conserved residues 3-11. . . . of peptide binding retains some of the std. features obsd. in MHC class Ia complexes, but novel features imply that HLA-E has evolved to mediate specific binding to a tightly defined set of almost identical hydrophobic peptides from the highly conserved class I leader sequences. These mol. adaptations make HLA-E a rigorous checkpoint at the cell surface reporting on the integrity of the antigen processing pathway to CD94/NKG2 receptor-bearing natural killer cells.
- IT CD antigens  
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(CD94, -NKG2 receptors; structural features impose tight peptide binding specificity in the nonclassical MHC mol. HLA-E and natural killer cell)
- IT Histocompatibility antigens  
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(HLA-E, complexes with peptides and CD94/NKG2 receptors; structural features impose tight peptide binding specificity in the nonclassical MHC mol. HLA-E)
- IT Antigen receptors  
RL: BPR (Biological process); BSU (Biological study, unclassified); PRP (Properties); BIOL (Biological study); PROC (Process)  
(NKG2/CD94 complex; structural features impose tight peptide binding specificity in the nonclassical MHC mol. HLA-E and natural killer cell)
- L8 ANSWER 12 OF 30 CAPLUS COPYRIGHT 2002 ACS  
SO Periodicum Biologorum (1998), 100(4), 441-443  
CODEN: PDBIAD; ISSN: 0031-5362
- IT Histocompatibility antigens  
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)  
(HLA-E; recognition by CD94/NKG2 heterodimers of human natural killer cells)
- L8 ANSWER 13 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 10  
TI HLA-E is the ligand for the natural killer cell CD94/NKG2 receptors  
SO Journal of Biomedical Science (Basel) (1998), 5(5), 321-331  
CODEN: JBCIEA; ISSN: 1021-7770
- AB A review is given with 102 refs. CD94/NKG2 is a recently described receptor present on natural killer (NK) cells and certain T cells that is composed of the CD94 chain covalently assocd. with a member of the NKG2 family of mols. Both chains are glycosylated members of the C-type lectin superfamily. The CD94/NKG2 receptors are functionally heterogeneous depending on which NKG2 family member is assocd. with CD94. It was thought that CD94/NKG2 receptors recognized a broad array of HLA-A, -B, and -C (classical), as well as the nonclassical HLA-G, MHC class I mols. Recent data have suggested that this receptor is specific for HLA-E complexed with a peptide derived from the signal sequence (residues 3-11) of certain classical MHC class I mols. Position 2 (residue 4) in the signal sequence derived peptides appears pivotal in detg. whether the HLA-E/peptide complex confers resistance to NK-mediated lysis. The potential roles that the CD94/NKG2-HLA-E receptor ligand interaction might play in infection and tumor development are discussed.
- IT CD antigens  
RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence)  
(CD94; HLA-E, ligand for the natural killer cell CD94/NKG2 receptors)
- IT Histocompatibility antigens  
RL: BAC (Biological activity or effector, except adverse); BOC (Biological occurrence); BSU (Biological study, unclassified); BIOL (Biological



study); OCCU (Occurrence)  
 (HLA; HLA-E, ligand for the natural  
 killer cell CD94/NKG2 receptors)

IT Proteins, specific or class  
 RL: BAC (Biological activity or effector, except adverse); BOC (Biological  
 occurrence); BSU (Biological study, unclassified); BIOL (Biological  
 study); OCCU (Occurrence)  
 (NKG2; HLA-E, ligand for the natural  
 killer cell CD94/NKG2 receptors)

IT Lymphocyte  
 (natural killer cell; HLA-E, ligand for the natural  
 killer cell CD94/NKG2 receptors)

L8 ANSWER 14 OF 30 CAPLUS COPYRIGHT 2002 ACS  
 SO Immunity (1998), 9(3), 289-294  
 CODEN: IUNIEH; ISSN: 1074-7613

AB A review with 50 refs. This paper discusses how recognition of  
 HLA-E enables NK cells to monitor the  
 integrity of the MHC class I-dependent antigen presentation pathway.  
 Since certain subpopulations of activated cytotoxic T cells (CTLs) can  
 also express the HLA-E-recognizing inhibitory  
 receptors, exptl. results are described regarding the effects of  
 inhibitory receptors on CTL activation. Finally, the potential  
 implications of HLA-E recognition on antitumor  
 immunity, antiviral immunity, and materno-fetal interactions are  
 discussed.

ST review cytotoxic lymphocyte HLA E CD94  
 NKG2

IT CD antigens  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
 (Biological study); PROC (Process)  
 (CD94; cytotoxic lymphocyte recognition of HLA-  
 E)

IT Receptors  
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL  
 (Biological study); PROC (Process)  
 (NKG2; cytotoxic lymphocyte recognition of HLA-  
 E)

L8 ANSWER 15 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 11  
 SO Immunological Reviews (1998), 163, 129-138  
 CODEN: IMRED2; ISSN: 0105-2896

AB A review with 49 refs. The major histocompatibility (MHC) class Ib mols.  
 HLA-E, HLA-F and HLA-G are relatively  
 non-polymorphic compared to class Ia mols. Both HLA-E  
 and HLA-G bind peptides and are involved in natural killer (  
 NK)-cell recognition, but the role of HLA-F is unclear.  
 HLA-E binds specifically to the conserved leader  
 sequence peptides from the class Ia MHC mols. and interacts on the cell  
 surface with the CD94/NKG2 class of NK-cell  
 receptors. The framework structure of HLA-E is  
 similar to that of the MHC class Ia mols., but the peptide-binding groove  
 is highly adapted for the specific. . . binding of the leader sequence  
 peptides. This is different from class Ia mols., which have highly  
 promiscuous peptide-binding grooves. The HLA-E groove  
 makes full use of all the available pockets and imposes specificity along  
 the entire length of the peptide. HLA-G. . . with leucine or  
 isoleucine at position 2, proline at position 3 and leucine at position 9.  
 Expression of HLA-G inhibits NK cells expressing the  
 CD94/NKG2 class of receptors, though an interaction with  
 these receptors has not been directly demonstrated.

IT CD antigens  
 RL: BSU (Biological study, unclassified); BIOL (Biological study)  
 (CD94; structure and function of human MHC class Ib mols.  
 HLA-E, HLA-F and HLA-G)

L8 ANSWER 16 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
 TI HLA-E-bound peptides influence recognition by  
 inhibitory and triggering CD94/NKG2 receptors:  
 Preferential response to an HLA-G-derived nonamer.

SO Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 80.  
 Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity  
 Seventeenth International. . .

IT Major Concepts  
 Immune System (Chemical Coordination and Homeostasis)

IT Chemicals & Biochemicals  
 CD94/NKG2 receptors; HLA-E  
 -bound peptides; HLA-G-derived nonamers

L8 ANSWER 17 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
 TI Decidual NK cells have receptors for HLA-E  
 which is expressed by human trophoblast.

SO Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 79.  
 Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity  
 Seventeenth International. . .

L8 ANSWER 18 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
 TI Specific recognition of HLA-E but not classical HLA  
 class I molecules by soluble CD94/NKG2A and NK  
 cells.

SO Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 72.  
 Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity  
 Seventeenth International. . .

IT . . .  
 Parts, Structures, & Systems of Organisms  
 NK cell [natural killer cell]: blood and lymphatics, immune system

IT Chemicals & Biochemicals  
 CD94; CD94/NKG2A complex; HLA class I  
 molecules; HLA-E: recognition; NKG2A

L8 ANSWER 19 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
 TI NK cell-mediated recognition of HLA-E and  
 HLA-G class Ib molecules.

SO Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 71.  
 Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity  
 Seventeenth International. . .

IT . . .  
 Parts, Structures, & Systems of Organisms  
 NK cell [natural killer cell]: blood and lymphatics, immune system

IT Chemicals & Biochemicals  
 CD94; CD94/NKG2A complex; HLA-  
 E class Ib molecules; HLA-G class Ib molecules; NKG2A

; SHP1 tyrosine phosphatase

- L8 ANSWER 20 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 12  
SO Tissue Antigens (1997), 50(6), 695-698  
CODEN: TSANA2; ISSN: 0001-2815
- AB . . . has also been found in both Cercopithecinae and humans. These allelic and evolutionary data suggest an altogether different functionality for HLA-E (and also HLA-G) compared with classical class I proteins: i.e., sending neg. (tolerogenic) signals to NK and T cells.
- L8 ANSWER 21 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 13  
SO Journal of Experimental Medicine (1997), 185(3), 385-391  
CODEN: JEMEDV; ISSN: 0022-1007
- AB . . . -B, and -C, is the presentation of peptides to T cells. A second function is the inhibition of natural killer (NK) cells, mediated by binding of class I mols. to NK receptors. In contrast, the function of the nonclassical human MHC class I mols., HLA-E, -F, and -G, is still a mystery. The specific expression of HLA-G in placental trophoblast suggests an important role for. . . escapes maternal allorecognition by downregulation of HLA-A and HLA-B mols. at this interface. It has been suggested that the maternal NK recognition of this downregulation is balanced by the expression of HLA-G, thus preventing damage to the placenta. Here, we describe the partial inhibition of NK lysis of the MHC class I neg. cell line LCL721.221 upon HLA-G transfection. We present three NK lines that are inhibited via the interaction of their NKAT3 receptor with HLA-G and with HLA-Bw4 mols. Inhibition can be blocked by the anti-NKAT3 antibody 5.133. In conclusion, NK inhibition by HLA-G via NKAT3 may contribute to the survival of the fetal semiallograft in the mother during pregnancy.
- L8 ANSWER 22 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
SO Placenta, (1997) Vol. 18, No. 2-3, pp. 234-235.  
ISSN: 0143-4004.
- IT Miscellaneous Descriptors  
BLOOD AND LYMPHATICS; CD4-POSITIVE CELL; CD8-POSITIVE CELL;  
CD94; HLA-DQ; HLA-DR; HLA-E; HLA-G; IMMUNE  
SYSTEM; INTERFERON-GAMMA; INTERLEUKIN-3; INTERLEUKIN-5; KILLER  
INHIBITORY RECEPTORS; LARGE GRANULAR LYMPHOCYTES; MAJOR  
HISTOCOMPATIBILITY COMPLEX; PEPTIDE INTERACTIONS; P58; RECURRENT  
SPONTANEOUS. . .
- L8 ANSWER 23 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 14  
SO Medecine/Sciences, (1996) 12/11 (1209-1218).  
ISSN: 0767-0974 CODEN: MSMSB4
- AB . . . encode membrane-anchored cell surface glycoproteins that present the endogenous antigenic peptides to the T cell receptors and are recognized by NK cell receptors. These genes are characterized by a high polymorphism and a nearly ubiquitous expression. The biological function of the three nonclassical class I genes HLA-E, HLA-F, and HLA-G still remains uncertain. In addition to these six genes, the MHC contains a number of class I pseudoalenes. . .
- L8 ANSWER 24 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 15  
SO J. Exp. Med. (1994), 180(2), 537-43  
CODEN: JEMEDV; ISSN: 0022-1007
- AB Natural killer (NK) cells kill normal and transformed hematopoietic cells that lack expression of major histocompatibility complex (MHC) class I antigens. Lysis of HLA-neg. Epstein Barr virus-transformed B lymphoblastoid cell lines (B-LCL) by human NK cell clones can be inhibited by transfection of the target cells with certain HLA-A, -B, or -C alleles. NK cell clones established from an individual demonstrate clonal heterogeneity in HLA recognition and a single NK clone can recognize multiple alleles. The authors describe a potential human NK cell receptor (NKB1) for certain HLA-B alleles (e.g., HLA-B\*5101 and -B\*5801) identified by the mAb DX9. NKB1 is a 70 kDa glycoprotein that is expressed on a subset of NK cells and NK cell clones. DX9 monoclonal antibody (mAb) specifically inhibits the interaction between NK cell clones and B-LCL targets transfected with certain HLA-B alleles, but does not affect recognition of HLA-A or HLA-C antigens. An individual NK cell clone can independently recognize B-LCL targets transfected with HLA-B or HLA-C antigens; however, DX9 mAb only affects interaction with transfectants expressing certain HLA-B alleles. These findings demonstrate the existence of NK cell receptors involved in the recognition of HLA-B and imply the presence of multiple receptors for MHC on an individual NK clone.
- L8 ANSWER 25 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 16  
SO Proc. Natl. Acad. Sci. U. S. A. (1992), 89(17), 8337-41  
CODEN: PNASAG; ISSN: 0027-8424
- AB Target structures important for natural killer (NK) cell recognition of virally infected cells are not well defined. Since major histocompatibility complex (MHC) class I mols. bind viral. . . acute infection, it was evaluated whether an interaction between MHC and virus might influence the susceptibility of infected cells to NK cell-mediated lysis. To control for MHC class I expression on target cells, either HLA class I-deficient C1R cells or C1R sublines expressing transfected HLA class I gene products were used. Human NK cells were unable to preferentially lyse class I-deficient C1R cells after infection with herpes simplex virus (HSV). In contrast, HLA class I transfectants were more susceptible to NK cell-mediated cytotoxicity after HSV infection. This occurred for HSV-infected C1R cells expressing any of the 3 HLA class I gene products tested (i.e., HLA-B27, HLA-A3, or HLA-Aw68), indicating that NK cells recognition in this system does not require self MHC and is not unique for a single haplotype. Productive HSV. . . the target structures on HSV-infected, HLA class I+ targets. These results demonstrate a role for MHC class I in regulating NK cell-mediated killing of virus-infected cells.
- L8 ANSWER 26 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 17  
SO Journal of Immunology, (1992) 148/2 (627-632).  
ISSN: 0022-1767 CODEN: JOIMA3
- AB . . . designated EO15 has been isolated and found to encode a non-classical HLA class I gene transcript. EO15 was compared with HLA-E and found to be 99.9% similar at the nucleotide level and to extend further in the 3' untranslated region. The. . . EO15 3' end suggests that EO15 clone represents a copy of the 3.3-kb mRNA species detected in Northern blot analyses. HLA-E transcripts of 1.9 and 3.3 kb have been described in a variety of cell

types. The two E015 mRNA species, similar in size to the previously defined HLA-E mRNA, were present at high levels in blood leukocyte populations and at variable levels in different cell lines. The E015. . . of E015 transcripts were detected in B and monocytic cell lines, whereas intermediate and lower levels were found in eosinophilic, NK-like, megakaryocytic, and T cell lines, respectively. Similar to its effect on classical HLA class I transcripts, IFN- $\gamma$  increased the levels. . . mRNA in eosinophils and neutrophils from hypereosinophilic patients. These results suggest that purified blood eosinophils as well as neutrophils express E015/HLA-E mRNA; however, further experiments are needed to investigate the localization and the function of E015 protein products.

L8 ANSWER 27 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.  
SO International Archives of Allergy and Immunology, (1992) 99/2-4  
(230-233).  
ISSN: 1018-2438 CODEN: IAAIEG

AB This paper reviews recent progresses on human Natural Killer (NK) cells which profoundly changed our concepts on NK cells and their functions. Regarding the ontogeny of NK cells, immature thymocyte populations (CD3-4-16-) have been shown to give rise to NK cells, provided suitable culture conditions. This indicates that precursor potentially capable of differentiation towards the NK cell lineage are present within the human thymus. Although NK cells lack known receptors for antigen (i.e. sIg and TCR), NK cell populations or clones were found to be capable of mediating specific recognition of allogenic cells. This specific function was clonally distributed and, more importantly, NK clones displaying different patterns of allospecificity could be isolated from single individuals. These data indicate the existence of a NK cell repertoire for alloantigen recognition. Analysis of the surface molecules identified by NK cells indicated that certain HLA alleles (e.g. HLA-Cw3) can act as specific protective elements from lysis by clones with defined specificities (e.g. specificity 2). Therefore, HLA class I molecules appear to play a central role in the NK cell-mediated functions. The finding that human NK cells express a clonally-distributed ability to recognize alloantigens suggested the existence of distinct surface receptors. Indeed, a new family of. . . the use of monoclonal antibodies. The expression of these molecules was shown to represent a stable phenotypic property of human NK cells and to be clonally distributed. More importantly, the expression of 58 kD molecules appears to correlate with the ability. . . given allospecificities. These data are supporting the concept that 58 kD molecules are part of receptor structures involved in the NK cell-mediated recognition.

L8 ANSWER 28 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 18  
SO Journal of Experimental Medicine, (1990) 171/4 (1315-1332).  
ISSN: 0022-1007 CODEN: JEMEA

AB. . . absence of CD4+ helper T cells and are not inhibited by anti-CD4 mAb. Both antigen-specific CTL as well as nonspecific NK cells can be elicited by dendritic cells. The NK cell response can be depleted at the precursor level by panning with an anti-CD11b mAb, which removes a CD11b+/CD28-, CD16+. . . have been sensitized by dendritic cells. The number of CD8+ cells that are blast transformed and express an activated phenotype (i.e., HLA DR/DQ+, CD25/IL-2R+, CD45R-) reaches 30-40% of the culture at day 4-5, the peak of the helper-independent response. We conclude that. . .

L8 ANSWER 29 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 19  
SO Journal of Clinical Investigation, (1989) 83/1 (278-287).  
ISSN: 0021-9738 CODEN: JCINAO

AB The role of major histocompatibility gene products (i.e., HLA molecules) in rendering tumor cells resistant to natural killer (NK) cell-mediated lysis was investigated by using mouse monoclonal antibodies to bind and mask HLA or non-HLA gene products on the. . . tumor targets. Enhanced lysis was not due to antibody dependent cellular cytotoxicity or due to an effect of antibody on NK effectors. Of importance, normal autologous and allogeneic human lymphocytes could not be lysed by NK cells despite blast transformation with mitogens or masking of HLA membrane determinants on blasts with monoclonal antibodies. Enhanced lysis, in the presence of antibody to HLA antigens, was not due to increased NK cell binding to tumor targets, but a consequence of enhanced postbinding lysis. Studies using granules obtained from NK cells indicated that masking of HLA antigens did not enhance the susceptibility of tumor targets to cytotoxicity. Such observations would. . . HLA antigens on tumor targets inhibit the triggering of effector cells (and release of cytotoxicins) after recognition and binding of NK cells to target cells.

L8 ANSWER 30 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 20  
SO Wiener Klinische Wochenschrift, (1987) 99/24 (848-855).  
ISSN: 0043-5325 CODEN: WKWOAO

AB. . . in conjunction with BAL provides further differentiation of lymphocytes into T- and B-, T-helper and T-suppressor types, natural killer cells (NK cells) and cytotoxic T-cells. A predominance of T-suppressor lymphocytes is an indication of EAA, whereas a predominance of T-helper lymphocytes is positively correlated with sarcoidosis. Other markers, e.g. HLA-DR, when expressed on the surface of alveolar macrophages, merely indicate activation unrelated to a specific type of lymphocytic alveolitis. BAL. . .

=> dis his

(FILE 'HOME' ENTERED AT 13:53:22 ON 24 JUN 2002)

FILE 'MEDLINE, CAPLUS, EMBASE, BIOSIS' ENTERED AT 13:53:34 ON 24 JUN 2002  
L1 3511 S BRAUD V?/AU OR ALLAN D?/AU OR OGG G?/AU OR OCALLAGHAN C?/AU O  
L2 47 S L1 AND (NK?)  
L3 35 S L2 AND (HLA (1N) E)  
L4 14 DUP REM L3 (21 DUPLICATES REMOVED)  
L5 72958 S NK (P) (NK? OR CD94?)  
L6 315 S (HLA (1N) E) (P) (NK? OR CD94?)  
L7 63 S L6 AND PD<19981204  
L8 30 DUP REM L7 (33 DUPLICATES REMOVED)

=> dis 18 1-30 ibib abs

L8 ANSWER 1 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 1  
ACCESSION NUMBER: 1998:289045 CAPLUS

DOCUMENT NUMBER: 129:66613  
 TITLE: HLA-E is a major ligand for the natural killer inhibitory receptor CD94/NKG2A

AUTHOR(S): Lee, Ni; Llano, Manuel; Carretero, Marta; Ishitani, Akiko; Navarro, Francisco; Lopez-Botet, Miguel; Geraghty, Daniel E.

CORPORATE SOURCE: The Clinical Research Division, Fred Hutchinson Cancer Research Center, Seattle, WA, 98104-2092, USA

SOURCE: Proceedings of the National Academy of Sciences of the United States of America (1998), 95(9), 5199-5204  
 CODEN: PNASA6; ISSN: 0027-8424

PUBLISHER: National Academy of Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The authors previously showed that the availability of a nonamer peptide derived from certain HLA class I signal sequences is a necessary requirement for the stabilization of endogenous HLA-E expression on the surface of 721.221 cells. This led the authors to examine the ability of HLA-E to protect HLA class I transfectants from natural killer (NK) cell-mediated lysis. It was possible to implicate the CD94/NKG2A complex as an inhibitory receptor recognizing this class Ib mol. by using as target a .221 transfectant selectively expressing surface HLA-E. HLA-E had no apparent inhibitory effect mediated through the identified Ig superfamily (Ig-SF) human killer cell inhibitory receptors or ILT2/LIR1. Further studies of CD94/NKG2+ NK cell-mediated recognition of .221 cells transfected with different HLA class I allotypes (i.e., -Cw4, -Cw3, -B7) confirmed that the inhibitory interaction was mediated by CD94/NKG2A recognizing the surface HLA-E mol., because only antibodies directed against either HLA-E, CD94, or CD94/NKG2A specifically restored lysis. Surface stabilization of HLA-E in cold-treated .221 cells loaded with appropriate peptides was sufficient to confer protection, resulting from recognition of the HLA class Ib mol. by the CD94/NKG2A inhibitory receptor. Consistent with the prediction that the ligand for CD94/NKG2A is expressed ubiquitously, the authors' examn. of HLA-E antigen distribution indicated that it is detectable on the surface of a wide variety of cell types.

L8 ANSWER 2 OF 30 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1998:288970 CAPLUS  
 DOCUMENT NUMBER: 129:66532  
 TITLE: HLA class I specificity for natural killer cell receptor CD94/NKG2A: two for one in more ways than one

AUTHOR(S): Yokoyama, Wayne M.

CORPORATE SOURCE: Howard Hughes Medical Institute, Rheumatology Division, Washington University School of Medicine, St. Louis, MO, 63110, USA

SOURCE: Proceedings of the National Academy of Sciences of the United States of America (1998), 95(9), 4791-4794  
 CODEN: PNASA6; ISSN: 0027-8424

PUBLISHER: National Academy of Sciences

DOCUMENT TYPE: Journal; General Review

LANGUAGE: English

AB A review and discussion with 44 refs. There are abundant data to substantiate the conclusion that CD94/NKG2A receptors of human natural killer cells directly recognize HLA-E mols.

L8 ANSWER 3 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 2  
 ACCESSION NUMBER: 1998:800568 CAPLUS  
 DOCUMENT NUMBER: 130:64947  
 TITLE: The Qa-1b molecule binds to a large subpopulation of murine NK cells

AUTHOR(S): Salcedo, Margarita; Bousso, Philippe; Ljunggren, Hans-Gustaf; Kourilsky, Philippe; Abastado, Jean-Pierre

CORPORATE SOURCE: Unite Biologie Moleculaire Gene, Institut Pasteur, Paris, F-75015, Fr.

SOURCE: European Journal of Immunology (1998), 28(12), 4356-4361  
 CODEN: EJIMAF; ISSN: 0014-2980

PUBLISHER: Wiley-VCH Verlag GmbH

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Recent studies on human NK cells have demonstrated that the NK cell CD94/NKG2 receptors bind to the nonclassical MHC class I mol. HLA-E. A functional CD94/NKG2 complex has not yet been identified in rodents, but cDNA encoding rat and mouse CD94 and NKG2 have recently been cloned, suggesting that CD94/NKG2 receptors may exist in species other than man. The mouse nonclassical MHC class I mol. Qa-1 shares several features with HLA-E. This suggests that Qa-1 may be similarly recognized by murine NK cells. To study the ability of Qa-1 to bind to murine NK cells, the authors have produced a sol. tetrameric form of Qa-1b. The authors demonstrate that Qa-1b tetramers distinctly bind to a large subset of fresh or IL-2-activated NK1.1+/CD3- splenocytes independently of the expression of Ly49 inhibitory receptors. Binding occurs whether NK cells have evolved in an MHC class I-expressing or in an MHC class I-deficient environment. The data suggest the existence of a Qa-1-recognizing structure on a large subpopulation of murine NK cells that may be similar to the human CD94/NKG2 heterodimeric complex.

REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 4 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 3  
 ACCESSION NUMBER: 1998:612269 CAPLUS  
 DOCUMENT NUMBER: 130:1528  
 TITLE: HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors. Preferential response to an HLA-G-derived nonamer

AUTHOR(S): Llano, Manuel; Lee, Ni; Navarro, Francisco; Garcia, Pilar; Albar, Juan Pablo; Geraghty, Daniel E.; Lopez-Botet, Miguel

CORPORATE SOURCE: Servicio Immunologia, Hospital Princesa, Madrid,  
E-28006, Spain  
SOURCE: European Journal of Immunology (1998),  
28(9), 2854-2863  
CODEN: EJIMAF; ISSN: 0014-2980  
PUBLISHER: Wiley-VCH Verlag GmbH  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The HLA-E class Ib mol. constitutes a major ligand for the lectin-like CD94/NKG2 natural killer (NK) cell receptors. Specific HLA class I leader sequence-derived nonapeptides bind to endogenous HLA-E mols. in the HLA-defective cell line 721.221, inducing HLA-E surface expression, and promote CD94/NKG2A-mediated recognition. The authors compared the ability of NK clones which expressed either inhibitory or activating CD94/NKG2 receptors to recognize HLA-E mols. on the surface of 721.221 cells loaded with a panel of synthetic nonamers derived from the leader sequences of most HLA class I mols. The results support the notion that the primary structure of the HLA-E-bound peptides influences CD94/NKG2-mediated recognition, beyond their ability to stabilize surface HLA-E. CD94/NKG2A+ NK clones appeared more sensitive to the interaction with most HLA-E-peptide complexes than did effector cells expressing the activating CD94/NKG2C receptor. An exception to this pattern was I-ILA-E loaded with the HLA-G-derived nonamer. This complex triggered cytotoxicity very efficiently over a wide range of peptide concns., suggesting that the HLA-E/G-nonamer complex interacts with the CD94/NKG2 triggering receptor with a higher affinity. These results raise the possibility that CD94/NKG2-mediated recognition of HLA-E expressed on extravillous cytotrophoblasts plays an important role in maternal-fetal cellular interactions.

REFERENCE COUNT: 47 THERE ARE 47 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 5 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 4

ACCESSION NUMBER: 1998:231442 CAPLUS

DOCUMENT NUMBER: 129:3692

TITLE: Specific engagement of the CD94/NKG2  
-A killer inhibitory receptor by the HLA-E class Ib molecule induces SHP-1 phosphatase recruitment to tyrosine-phosphorylated NKG2  
-A. Evidence for receptor function in heterologous transfectants

AUTHOR(S): Carretero, Marta; Palmieri, Gabriella; Llano, Manuel; Tullio, Valentino; Santoni, Angela; Geraghty, Daniel E.; Lopez-Botet, Miguel

CORPORATE SOURCE: Servicio Immunologia, Hospital Princesa, Universidad Autonoma Madrid, Madrid, E-28006, Spain

SOURCE: European Journal of Immunology (1998),  
28(4), 1280-1291

CODEN: EJIMAF; ISSN: 0014-2980

PUBLISHER: Wiley-VCH Verlag GmbH

DOCUMENT TYPE: Journal

LANGUAGE: English

AB It was recently demonstrated that the CD94/NKG2-A killer inhibitory receptor (KIR) specifically recognizes the HLA-E class Ib mol. The apparent CD94-mediated specific recognition of different HLA class Ia allotypes, transfected into the HLA-defective cell line 721.221, indeed depends on their selective ability to concomitantly stabilize the surface expression of endogenous HLA-E mols., which confer protection against CD94/NKG2-A+ effector cells. The authors show that a selective engagement of the CD94/NKG2-A inhibitory receptor with a specific monoclonal antibody (mAb) (Z199) was sufficient to induce Tyr phosphorylation of the NKG2-A subunit and SHP-1 recruitment. These early biochem. events, commonly related to neg. signaling pathways, were also detected upon the specific interaction of NK cells with an HLA-E+ 721.221 transfectant (.221-AEH), and were prevented by pre-incubation of .221-AEH with an anti-HLA class I mAb. MAb crosslinking of the CD94/NKG2-A receptor, segregated from other NK-assocd. mols. by transfection into a rat basophilic leukemia cell line (RBL-2H3), promoted Tyr phosphorylation of NKG2-A and co-pptn. of SHP-1, together with an inhibition of secretory events triggered via Fc.epsilon.RI. Interaction of CD94/NKG2-A+ RBL cells with the HLA-E+.221-AEH transfectant specifically induced a detectable assocn. of SHP-1 with NKG2-A, constituting a more formal evidence for the receptor-HLA class I interaction.

L8 ANSWER 6 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 5

ACCESSION NUMBER: 1998:602367 CAPLUS

DOCUMENT NUMBER: 129:301413

TITLE: Qa-1b binds conserved class I leader peptides derived from several mammalian species

AUTHOR(S): Kurepa, Zoran; Hasemann, Charles A.; Forman, James  
CORPORATE SOURCE: Immunology Graduate Program, University of Texas Southwestern Medical Center at Dallas, Dallas, TX, 75235, USA

SOURCE: Journal of Experimental Medicine (1998),  
188(5), 973-978

CODEN: JEMEAU; ISSN: 0022-1007

PUBLISHER: Rockefeller University Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Qa-1b binds a peptide (AMAPRTLLL), referred to as Qdm (for Qa-1 determinant modifier), derived from the signal sequence of murine class Ia mols. This peptide binds with high affinity and accounts for almost all of the peptides assocd. with this mol. Human histocompatibility leukocyte antigen (HLA)-E, a homolog of Qa-1b, binds similar peptides derived from human class Ia mols. and interacts with CD94/NKG2 receptors on natural killer cells. The authors used surface plasmon resonance to det. the ability of Qa-1b to bind related ligands representing peptides derived from the leaders of class I mols. from several mammalian species. All of the peptides reported to bind HLA-E bound readily to Qa-1b. In addn., peptides derived from leader segments of different mammals also bound to Qa-1b, indicating a conservation of this "Qdm-like" epitope throughout mammalian evolution. The authors have attempted to define a minimal peptide on a polyglycine backbone that binds Qa-1b. Previous findings showed that P2

and P9 are important but not sufficient for binding to Qa-1b. Although a min. peptide (GMGGGGLLL) bound Qa-1b, its interaction was relatively weak, as were peptides sharing five or six residues with Qdm, indicating that multiple native residues are required for a strong interaction. This finding is consistent with the observation that this mol. preferentially binds this single ligand.

L8 ANSWER 7 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 6  
 ACCESSION NUMBER: 1998:162486 CAPLUS  
 DOCUMENT NUMBER: 128:281696  
 TITLE: Recognition of human histocompatibility leukocyte antigen (HLA)-E complexed with HLA class I signal sequence-derived peptides by CD94/NKG2 confers protection from natural killer cell-mediated lysis  
 AUTHOR(S): Borrego, Francisco; Ulbrecht, Matthias; Weiss, Elisabeth H.; Coligan, John E.; Brooks, Andrew G.  
 CORPORATE SOURCE: Laboratory of Molecular Structure, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Rockville, MD, 20852, USA  
 SOURCE: Journal of Experimental Medicine (1998), 187(5), 813-818  
 CODEN: JEMEA; ISSN: 0022-1007  
 PUBLISHER: Rockefeller University Press  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Human histocompatibility leukocyte antigen (HLA)-E is a nonclassical HLA class I mol., the gene for which is transcribed in most tissues. It has recently been reported that this mol. binds peptides derived from the signal sequence of HLA class I proteins; however, no function for HLA-E has yet been described. The authors show that natural killer (NK) cells can recognize target cells expressing HLA-E mols. on the cell surface and this interaction results in inhibition of the lytic process. Furthermore, HLA-E recognition is mediated primarily through the CD94/NKG2-A heterodimer, as CD94-specific, but not killer cell inhibitory receptor (KIR)-specific mAbs block HLA-E-mediated protection of target cells. Cell surface HLA-E could be increased by incubation with synthetic peptides corresponding to residues 3-11 from the signal sequences of a no. of HLA class I mols.; however, only peptides which contained a Met at position 2 were capable of conferring resistance to NK-mediated lysis, whereas those having Thr at position 2 had no effect. Interestingly, HLA class I mols. previously correlated with CD94/NKG2 recognition all have Met at residue 4 of the signal sequence (position 2 of the HLA-E binding peptide), whereas those which have been reported not to interact with CD94/NKG2 have Thr at this position. These data thus show a function for HLA-E and suggest an alternative explanation for the apparent broad reactivity of CD94/NKG2 with HLA class I mols.; that CD94/NKG2 interacts with HLA-E complexed with signal sequence peptides derived from "protective" HLA class I alleles rather than directly interacting with classical HLA class I proteins.

L8 ANSWER 8 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 7  
 ACCESSION NUMBER: 1998:134074 CAPLUS  
 DOCUMENT NUMBER: 128:269312  
 TITLE: HLA-E binds to natural killer cell receptors CD94/NKG2A, B and C  
 AUTHOR(S): Braud, Veronique M.; Allan, David S. J.; O'Callaghan, Christopher A.; Soderstrom, Kalle; D'Andrea, Annalisa; Ogg, Graham S.; Lazatic, Sasha; Young, Neil T.; Bell, John I.; Phillips, Joseph H.; Lanier, Lewis L.; McMichael, Andrew J.  
 CORPORATE SOURCE: Inst. Molecular Med., John Radcliffe Hosp., Oxford, OX3 9DS, UK  
 SOURCE: Nature (London) (1998), 391(6669), 795-799  
 CODEN: NATUAS; ISSN: 0028-0836  
 PUBLISHER: Macmillan Magazines  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The protein HLA-E is a non-classical major histocompatibility complex (MHC) mol. of limited sequence variability. Its expression on the cell surface is regulated by the binding of peptides derived from the signal sequence of some other MHC class I mols. Here we report the identification of ligands for HLA-E. We constructed tetramers in which recombinant HLA-E and .beta.2-microglobulin were refolded with an MHC, leader-sequence peptide, biotinylated, and conjugated to phycoerythrin-labeled Extravidin. This HLA-E tetramer bound to natural killer (NK) cells and a small subset of T cells from peripheral blood. On transfectants, the tetramer bound to the CD94/NKG2A, CD94/NKG2B and CD94/NKG2C NM cell receptors, but did not bind to the Ig family of NK cell receptors (KIR). Surface expression of HLA-E was enough to protect target cells from lysis by CD94/NKG2A + NK-cell clones. A subset of HLA class I alleles has been shown to inhibit killing by CD94/NKG2A+ NK-cell clones. Only the HLA alleles that possess a leader peptide capable of upregulating HLA-E surface expression confer resistance to NK-cell-mediated lysis, implying that their action is mediated by HLA-E, the predominant ligand for the NK cell inhibitory receptor CD94/NKG2A.

L8 ANSWER 9 OF 30 CAPLUS COPYRIGHT 2002 ACS  
 ACCESSION NUMBER: 1998:226619 CAPLUS  
 DOCUMENT NUMBER: 128:320229  
 TITLE: Follow the leader: NK cell receptors for classical and nonclassical MHC class I  
 AUTHOR(S): Lanier, Lewis L.  
 CORPORATE SOURCE: Immunobiology Department, DNAX Research Institute of Molecular and Cellular Biology, Palo Alto, CA, 94304, USA  
 SOURCE: Cell (Cambridge, Massachusetts) (1998), 92(6), 705-707  
 CODEN: CELLB5; ISSN: 0092-8674  
 PUBLISHER: Cell Press  
 DOCUMENT TYPE: Journal; General Review  
 LANGUAGE: English

AB A review, with 15 refs., discussing the function of mouse Ly49 receptors, the structure and function of human KIR receptors, and the mol.

interaction of HLA-E with CD94/NKG2 heterodimers.

L8 ANSWER 10 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 8  
ACCESSION NUMBER: 1998:417529 CAPLUS  
DOCUMENT NUMBER: 129:160384  
TITLE: Association of DAP12 with activating CD94/NKG2C NK cell receptors  
AUTHOR(S): Lanier, Lewis L.; Corliss, Brian; Wu, Jun; Phillips, Joseph H.  
CORPORATE SOURCE: Immunobiology Dept., DNAX Research Inst. of Molecular and Cellular Biology, Palo Alto, CA, 94304, USA  
SOURCE: Immunity (1998), 8(6), 693-701  
CODEN: IUMIEH; ISSN: 1074-7613  
PUBLISHER: Cell Press  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB While the inhibitory NK cell receptors for MHC class I express immunoreceptor tyrosine-based inhibitory motifs that recruit intracellular tyrosine phosphatases and prevent NK cell effector function, the activating NK cell receptors lack intrinsic sequences required for cellular stimulation. CD94/NKG2C, an activating NK cell receptor of the C-type lectin superfamily that binds to HLA-E, noncovalently assoc. with DAP12, a membrane receptor contg. an immunoreceptor tyrosine-based activating motif. Efficient expression of CD94/NKG2C on the cell surface requires the presence of DAP12, and charged residues in the transmembrane domains of DAP12 and NKG2C are necessary for this interaction. These results provide a mol. basis for the assembly of NK cell receptors for MHC class I involved in cellular activation and inhibition.

L8 ANSWER 11 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 9  
ACCESSION NUMBER: 1998:225128 CAPLUS  
DOCUMENT NUMBER: 129:3675  
TITLE: Structural features impose tight peptide binding specificity in the nonclassical MHC molecule HLA-E  
AUTHOR(S): O'callaghan, Christopher A.; Tormo, Jose; Willcox, Benjamin E.; Braud, Veronique M.; Jakobsen, Bent K.; Stuart, David I.; McMichael, Andrew J.; Bell, John I.; Jones, E. Yvonne  
CORPORATE SOURCE: Molecular Immunology Group, Nuffield Department of Clinical Medicine, Institute of Molecular Medicine, John Radcliffe Hospital, University of Oxford, Oxford, OX3 9DS, UK  
SOURCE: Molecular Cell (1998), 1(4), 531-541  
CODEN: MOCEFL; ISSN: 1097-2765  
PUBLISHER: Cell Press  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The crystal structure of the nonclassical human class Ib MHC mol. HLA-E has been detd. in complex with a prototypic ligand, the nonamer peptide (VMAPRTVLL), derived from the highly conserved residues 3-11 of the human MHC class Ia leader sequence. The mode of peptide binding retains some of the std. features obsd. in MHC class Ia complexes, but novel features imply that HLA-E has evolved to mediate specific binding to a tightly defined set of almost identical hydrophobic peptides from the highly conserved class I leader sequences. These mol. adaptations make HLA-E a rigorous checkpoint at the cell surface reporting on the integrity of the antigen processing pathway to CD94/NKG2 receptor-bearing natural killer cells.

L8 ANSWER 12 OF 30 CAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 1999:179237 CAPLUS  
DOCUMENT NUMBER: 131:17585  
TITLE: Human C-type lectin receptors involved in NK cell mediated recognition of HLA class I molecules  
AUTHOR(S): Bellon, Teresa; Navarro, Francisco; Llano, Manuel; Garcia, Pilar; Lopez-Botet, Miguel  
CORPORATE SOURCE: Servicio de Inmunologia, Hospital Universitario de la Princesa, Madrid, 28006, Spain  
SOURCE: Periodicum Biologorum (1998), 100(4), 441-443  
CODEN: PDBIAD; ISSN: 0031-5362  
PUBLISHER: Hrvatsko Prirodoslovno Društvo  
DOCUMENT TYPE: Journal; General Review  
LANGUAGE: English  
AB A review with 42 refs. discussing the function of CD94/NKG2 heterodimers and their ligand specificity.  
REFERENCE COUNT: 42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 13 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 10  
ACCESSION NUMBER: 1998:685732 CAPLUS  
DOCUMENT NUMBER: 129:288814  
TITLE: HLA-E is the ligand for the natural killer cell CD94/NKG2 receptors  
AUTHOR(S): Posch, Phillip E.; Borrego, Francisco; Brooks, Andrew G.; Coligan, John E.  
CORPORATE SOURCE: Structural Biology Section, National Inst. Allergy Infectious Disease, National Inst. Health, Rockville, MD, 20852, USA  
SOURCE: Journal of Biomedical Science (Basel) (1998), 5(5), 321-331  
CODEN: JBCIEA; ISSN: 1021-7770  
PUBLISHER: S. Karger AG  
DOCUMENT TYPE: Journal; General Review  
LANGUAGE: English  
AB A review is given with 102 refs. CD94/NKG2 is a recently described receptor present on natural killer (NK) cells and certain T cells that is composed of the CD94 chain covalently assocd. with a member of the NKG2 family of mols. Both chains are glycosylated members of the C-type lectin superfamily. The CD94/NKG2 receptors are functionally heterogeneous depending on which NKG2 family member is assocd. with CD94. It was thought that CD94/NKG2 receptors recognized a broad array of HLA-A, -B, and -C (classical), as well as the nonclassical HLA-G, MHC class I mols. Recent data have suggested that this receptor is specific for HLA-E complexed with a peptide derived from the signal sequence (residues 3-11) of certain classical MHC class I mols. Position 2 (residue 4) in the signal sequence

derived peptides appears pivotal in detg. whether the HLA-E/peptide complex confers resistance to NK-mediated lysis. The potential roles that the CD94/NKG2-HLA-E receptor ligand interaction might play in infection and tumor development are discussed.

L8 ANSWER 14 OF 30 CAPLUS COPYRIGHT 2002 ACS  
ACCESSION NUMBER: 1998:676185 CAPLUS  
DOCUMENT NUMBER: 130:36956  
TITLE: Cytotoxic lymphocyte recognition of HLA-E: utilizing a nonclassical window to peer into classical MHC  
AUTHOR(S): Leibson, Paul J.  
CORPORATE SOURCE: Department of Immunology, Mayo Clinic and Foundation, Rochester, MN, 55905, USA  
SOURCE: Immunity (1998), 9(3), 289-294  
CODEN: IUNIEH; ISSN: 1074-7613  
PUBLISHER: Cell Press  
DOCUMENT TYPE: Journal; General Review  
LANGUAGE: English  
AB A review with 50 refs. This paper discusses how recognition of HLA-E enables NK cells to monitor the integrity of the MHC class I-dependent antigen presentation pathway. Since certain subpopulations of activated cytotoxic T cells (CTLs) can also express the HLA-E-recognizing inhibitory receptors, exptl. results are described regarding the effects of inhibitory receptors on CTL activation. Finally, the potential implications of HLA-E recognition on antitumor immunity, antiviral immunity, and materno-fetal interactions are discussed.  
REFERENCE COUNT: 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 15 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 11  
ACCESSION NUMBER: 1998:543874 CAPLUS  
DOCUMENT NUMBER: 129:274318  
TITLE: Structure and function of the human MHC class Ib molecules HLA-E, HLA-F and HLA-G  
AUTHOR(S): O'Callaghan, Christopher A.; Bell, John I.  
CORPORATE SOURCE: Molecular Immunology Group, Nuffield Department Clinical Medicine, Institute Molecular Medicine, John Radcliffe Hospital, University Oxford, Oxford, UK  
SOURCE: Immunological Reviews (1998), 163, 129-138  
CODEN: IMRED2; ISSN: 0105-2896  
PUBLISHER: Munksgaard International Publishers Ltd.  
DOCUMENT TYPE: Journal; General Review  
LANGUAGE: English  
AB A review with 49 refs. The major histocompatibility (MHC) class Ib mols. HLA-E, HLA-F and HLA-G are relatively non-polymorphic compared to class Ia mols. Both HLA-E and HLA-G bind peptides and are involved in natural killer (NK)-cell recognition, but the role of HLA-F is unclear. HLA-E binds specifically to the conserved leader sequence peptides from the class Ia MHC mols. and interacts on the cell surface with the CD94/NKG2 class of NK-cell receptors. The framework structure of HLA-E is similar to that of the MHC class Ia mols., but the peptide-binding groove is highly adapted for the specific binding of the leader sequence peptides. This is different from class Ia mols., which have highly promiscuous peptide-binding grooves. The HLA-E groove makes full use of all the available pockets and imposes specificity along the entire length of the peptide. HLA-G binds nonamer peptides with leucine or isoleucine at position 2, proline at position 3 and leucine at position 9. Expression of HLA-G inhibits NK cells expressing the CD94/NKG2 class of receptors, though an interaction with these receptors has not been directly demonstrated.

L8 ANSWER 16 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
ACCESSION NUMBER: 1999:219822 BIOSIS  
DOCUMENT NUMBER: PREV199900219822  
TITLE: HLA-E-bound peptides influence recognition by inhibitory and triggering CD94/NKG2 receptors: Preferential response to an HLA-G-derived nonamer.  
AUTHOR(S): Llano, Manuel (1); Lee, Ni; Navarro, Francisco (1); Garcia, Pilar (1); Albar, Juan Pablo; Geraghty, Daniel E.; Lopez-Botet, Miguel (1)  
CORPORATE SOURCE: (1) Hospital Universitario de la Princesa, Madrid Spain  
SOURCE: Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 80.  
Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity Seventeenth International Natural Killer Cell Workshop Warrenton, Virginia, USA October 17-21, 1998  
ISSN: 1018-8916.  
DOCUMENT TYPE: Conference  
LANGUAGE: English

L8 ANSWER 17 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
ACCESSION NUMBER: 1999:219821 BIOSIS  
DOCUMENT NUMBER: PREV199900219821  
TITLE: Decidual NK cells have receptors for HLA-E which is expressed by human trophoblast.  
AUTHOR(S): Allan, D.S.J. (1); Verma, S.; Bowen, J. M.; Loke, Y. W.; McMichael, J. (1); Braud, V. M. (1); King, A.  
CORPORATE SOURCE: (1) Institute of Molecular Medicine, John Radcliffe Hospital, Oxford, OX3 9DS UK  
SOURCE: Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 79.  
Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity Seventeenth International Natural Killer Cell Workshop Warrenton, Virginia, USA October 17-21, 1998  
ISSN: 1018-8916.  
DOCUMENT TYPE: Conference  
LANGUAGE: English

L8 ANSWER 18 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
ACCESSION NUMBER: 1999:219811 BIOSIS  
DOCUMENT NUMBER: PREV199900219811  
TITLE: Specific recognition of HLA-E but not classical HLA class I molecules by soluble CD94/NKG2A and NK cells.  
AUTHOR(S): Brooks, Andrew G. (1); Borrego, Francisco; Posch, Phillip E. (1); Patamawenu, Apisit (1); Coligan, John E.



CORPORATE SOURCE: (1) Structural Biology Section, National Institute of Allergy and Infectious Disease, National Institutes of Health, Twinbrook II, Rockville, MD, 20852 USA  
SOURCE: Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 72.  
Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity Seventeenth International Natural Killer Cell Workshop Warrenton, Virginia, USA October 17-21, 1998  
ISSN: 1018-8916.  
DOCUMENT TYPE: Conference  
LANGUAGE: English

L8 ANSWER 19 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
ACCESSION NUMBER: 1999:219810 BIOSIS  
DOCUMENT NUMBER: PREV199900219810  
TITLE: NK cell-mediated recognition of HLA-E and HLA-G class Ib molecules.  
AUTHOR(S): Lopez-Botet, M. (1); Llano, M. (1); Navarro, F. (1); Garcia, P. (1)  
CORPORATE SOURCE: (1) S. de Immunologia, Hospital Universitario de la Princesa, 28006, Madrid Spain  
SOURCE: Natural Immunity, (Feb., 1998) Vol. 16, No. 2-3, pp. 71.  
Meeting Info.: Fifth Annual Meeting of the Society for Natural Immunity Seventeenth International Natural Killer Cell Workshop Warrenton, Virginia, USA October 17-21, 1998  
ISSN: 1018-8916.  
DOCUMENT TYPE: Conference  
LANGUAGE: English

L8 ANSWER 20 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 12  
ACCESSION NUMBER: 1998:23376 CAPLUS  
DOCUMENT NUMBER: 128:137001  
TITLE: Mhc-E polymorphism in Pongidae primates: the same allele is found in two different species  
AUTHOR(S): Suarez, B.; Morales, P.; Castro, M. J.; Fernandez-Soria, V.; Recio, M. J.; Perez-Blas, M.; Alvarez, M.; Diaz-Campos, N.; Arnaiz-Villena, A.  
CORPORATE SOURCE: Department of Immunology, Hospital 12 de Octubre, Universidad Complutense, Madrid, Spain  
SOURCE: Tissue Antigens (1997), 50(6), 695-698  
CODEN: TSANA2; ISSN: 0001-2815  
PUBLISHER: Munksgaard International Publishers Ltd.  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Mhc-E intron 1, exon 2, intron 2, and exon 3 from pygmy chimpanzee (*Pan paniscus*), chimpanzee (*Pan troglodytes*), gorilla (*Gorilla gorilla*) and orangutan (*Pongo pygmaeus*) have been sequenced; six new Mhc-E alleles have been obtained but sequence changes are only placed either in introns or in synonymous exonic bases. One pygmy chimpanzee Mhc-E DNA sequence is identical to another sequence from chimpanzee; the fact that no variation is found also at the intronic level suggests that these two species of chimpanzee may have recently sepd. and/or that both of them might only represent subspecies. Mhc-E phylogenetic trees sep. two evolutionary groups: Pongidae, including humans, and Cercopithecinae; this is also found by studying another non-classical class I gene, Mhc-G. The Mhc-E alleles' invariance at the protein level supports that strong selective forces are operating at the Mhc-E locus, as has also been found in both Cercopithecinae and humans. These allelic and evolutionary data suggest an altogether different functionality for HLA-E (and also HLA-G) compared with classical class I proteins: i.e., sending neg. (tolerogenic) signals to NK and T cells.

L8 ANSWER 21 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 13  
ACCESSION NUMBER: 1997:111333 CAPLUS  
DOCUMENT NUMBER: 126:184797  
TITLE: Human histocompatibility leukocyte antigen (HLA)-G molecules inhibit NKAT3 expressing natural killer cells  
AUTHOR(S): Muenz, Christian; Holmes, Nicholas; King, Ashley; Loke, Yung Wai; Colonna, Marco; Schild, Hansjoerg; Rammensee, Hans-Georg  
CORPORATE SOURCE: Dep. Immunology, Univ. Tuebingen, Tuebingen, 72076, Germany  
SOURCE: Journal of Experimental Medicine (1997), 185(3), 385-391  
CODEN: JEMEAU; ISSN: 0022-1007  
PUBLISHER: Rockefeller University Press  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The crucial immunol. function of the classical human major histocompatibility complex (MHC) class I mols., human histocompatibility leukocyte antigen (HLA)-A, -B, and -C, is the presentation of peptides to T cells. A second function is the inhibition of natural killer (NK) cells, mediated by binding of class I mols. to NK receptors. In contrast, the function of the nonclassical human MHC class I mols., HLA-E, -F, and -G, is still a mystery. The specific expression of HLA-G in placental trophoblast suggests an important role for this mol. in the immunol. interaction between mother and child. The fetus, semiallograft by its genotype, escapes maternal allorecognition by downregulation of HLA-A and HLA-B mols. at this interface. It has been suggested that the maternal NK recognition of this downregulation is balanced by the expression of HLA-G, thus preventing damage to the placenta. Here, we describe the partial inhibition of NK lysis of the MHC class I neg. cell line LCL721.221 upon HLA-G transfection. We present three NK lines that are inhibited via the interaction of their NKAT3 receptor with HLA-G and with HLA-Bw4 mols. Inhibition can be blocked by the anti-NKAT3 antibody 5.133. In conclusion, NK inhibition by HLA-G via NKAT3 may contribute to the survival of the fetal semiallograft in the mother during pregnancy.

L8 ANSWER 22 OF 30 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
ACCESSION NUMBER: 1997:227193 BIOSIS  
DOCUMENT NUMBER: PREV199799518909  
TITLE: The Thomas G. Wegmann Memorial Symposium on reproductive immunology: Banff, Alberta, Canada, (September 12, 1996).  
AUTHOR(S): Vince, G.  
CORPORATE SOURCE: Dep. Immunol., Univ. Liverpool, Duncan Build., Liverpool L69 3BX UK  
SOURCE: Placenta, (1997) Vol. 18, No. 2-3, pp. 234-235.  
ISSN: 0143-4004.

DOCUMENT TYPE: Conference; Report  
LANGUAGE: English

L8 ANSWER 23 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 14  
ACCESSION NUMBER: 96353415 EMBASE  
DOCUMENT NUMBER: 1996353415  
TITLE:

[The class I region of the MHC genes is one of the most complex in the whole human genome].  
LA REGION HLA DE CLASSE I: UNE ORGANISATION COMPLIQUEE PAR LA PRESENCE DE NOMBREUSES FAMILLES MULTIGENIQUES.  
Pichon L.; Giffon T.; Chauvel B.; Le Gall J.-Y.; David V.  
CORPORATE SOURCE: UPR Cnrs, Recombinations genetiques, Faculte de Medecine, 2 avenue du Pr-Leon Bernard, 35043 Rennes Cedex, France  
SOURCE: Medecine/Sciences, (1996) 12/11 (1209-1218).  
ISSN: 0767-0974 CODEN: MSMSE4  
COUNTRY: France  
DOCUMENT TYPE: Journal; General Review  
FILE SEGMENT: 022 Human Genetics  
026 Immunology, Serology and Transplantation  
029 Clinical Biochemistry

LANGUAGE: French  
SUMMARY LANGUAGE: French; English

AB The human major histocompatibility complex (MHC) is a 4-Mb genomic segment located on chromosome 6, in the distal portion of the 6p21.3 band. The MHC is a well described region of the human genome and represents a cluster of genes intensively studied because of the important functions of the gene products in the immune system. The total number of genes located within the MHC is now approximately 100, but more can be predicted. The density of genes in the Class II and Class III regions has been known for several years, but the high density of genes in the Class I region did only recently become apparent. This class I region spans 2,000 kb on the telomeric half of the MHC, and contains six class I genes, three of which, the classical human leukocyte antigen genes HLA-A, HLA-B and HLA-C, encode membrane-anchored cell surface glycoproteins that present the endogenous antigenic peptides to the T cell receptors and are recognized by NK cell receptors. These genes are characterized by a high polymorphism and a nearly ubiquitous expression. The biological function of the three nonclassical class I genes HLA-E, HLA-F, and HLA-G still remains uncertain. In addition to these six genes, the MHC contains a number of class I pseudogenes and gene fragments whose close relationship indicates a common origin from a typical HLA class I ancestor gene. This region appears also to harbour various non-HLA single copy genes such as OTF3, TCF19, S gene, TUBB, HSRL, B30.2, ZNF173, HCG V, HCG-VII, MOG or OLFR2 as well as eight recently described multigene families named P5.1, 3.8-1, PERB11, BAT1, HCGII, HCGIV, HCGVIII and HCGIX containing a number of sequences displayed throughout the HLA Class I region. The presence of all these coding sequences and multigene families suggests that the gene density of the HLA Class I region is comparable to that of the class II and class III regions, i.e. one gene per 20 or 30 kb. We present here a new map compiling the update knowledge of the structure of this Class I region.

L8 ANSWER 24 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 15

ACCESSION NUMBER: 1994:506185 CAPLUS  
DOCUMENT NUMBER: 121:106185  
TITLE: NKb1: a natural killer cell receptor involved in the recognition of polymorphic HLA-B molecules  
AUTHOR(S): Litwin, Virginia; Gumperz, Jenny; Parham, Peter; Phillips, Joseph H.; Lanier, Lewis L.  
CORPORATE SOURCE: Department of Human Immunology, DNAX Research Institute of Molecular Cellular Biology, Inc., Palo Alto, CA, 94304, USA  
SOURCE: J. Exp. Med. (1994), 180(2), 537-43  
CODEN: JEMEA; ISSN: 0022-1007  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Natural killer (NK) cells kill normal and transformed hematopoietic cells that lack expression of major histocompatibility complex (MHC) class I antigens. Lysis of HLA-neg. Epstein Barr virus-transformed B lymphoblastoid cell lines (B-LCL) by human NK cell clones can be inhibited by transfection of the target cells with certain HLA-A, -B, or -C alleles. NK cell clones established from an individual demonstrate clonal heterogeneity in HLA recognition and a single NK clone can recognize multiple alleles. The authors describe a potential human NK cell receptor (NKb1) for certain HLA-B alleles (e.g., HLA-B\*5101 and -B\*5801) identified by the mAb DX9. NKb1 is a 70 kDa glycoprotein that is expressed on a subset of NK cells and NK cell clones. DX9 monoclonal antibody (mAb) specifically inhibits the interaction between NK cell clones and B-LCL targets transfected with certain HLA-B alleles, but does not affect recognition of HLA-A or HLA-C antigens. An individual NK cell clone can independently recognize B-LCL targets transfected with HLA-B or HLA-C antigens; however, DX9 mAb only affects interaction with transfectants expressing certain HLA-B alleles. These findings demonstrate the existence of NK cell receptors involved in the recognition of HLA-B and imply the presence of multiple receptors for MHC on an individual NK clone.

L8 ANSWER 25 OF 30 CAPLUS COPYRIGHT 2002 ACS DUPLICATE 16

ACCESSION NUMBER: 1993:20561 CAPLUS  
DOCUMENT NUMBER: 118:20561  
TITLE: Role for major histocompatibility complex class I in regulating natural killer cell-mediated killing of virus-infected cells  
AUTHOR(S): Kaufman, Dan S.; Schoon, Renee A.; Leibson, Paul J.  
CORPORATE SOURCE: Dep. Immunol., Mayo Clin. Found., Rochester, MN, 55905, USA  
SOURCE: Proc. Natl. Acad. Sci. U. S. A. (1992), 89(17), 8337-41  
CODEN: PNASAG; ISSN: 0027-8424  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Target structures important for natural killer (NK) cell recognition of virally infected cells are not well defined. Since major histocompatibility complex (MHC) class I mols. bind viral peptides during acute infection, it was evaluated whether an interaction between MHC and virus might influence the susceptibility of infected cells to NK cell-mediated lysis. To control for MHC class I expression on target cells, either HLA class I-deficient C1R cells or C1R sublines expressing transfected HLA class I gene products were used. Human NK cells were unable to preferentially lyse class I-deficient C1R cells after infection with herpes simplex virus (HSV). In contrast, HLA class I

transfectants were more susceptible to NK cell-mediated cytotoxicity after HSV infection. This occurred for HSV-infected C1R cells expressing any of the 3 HLA class I gene products tested (i.e., HLA-B27, HLA-A3, or HLA-Aw68), indicating that NK cells recognition in this system does not require self MHC and is not unique for a single haplotype. Productive HSV infection is required for the increased killing, since inoculation with UV-inactivated virus did not lead to increased lysis. In addn., since HSV infection of the transfectants did not alter the level of class I expression, the change in susceptibility appears to be due to qual. changes in the target structures on HSV-infected, HLA class I+ targets. These results demonstrate a role for MHC class I in regulating NK cell-mediated killing of virus-infected cells.

L8 ANSWER 26 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 17  
 ACCESSION NUMBER: 92041666 EMBASE  
 DOCUMENT NUMBER: 1992041666  
 TITLE: Cloning and expression of a cDNA encoding a non-classical MHC class I antigen (HLA-E) in eosinophils from hypereosinophilic patients.  
 AUTHOR: Truong M.-J.; Gruart V.; Capron A.; Capron M.; Tourville B.  
 CORPORATE SOURCE: Ctr. d'Immunol. Biol. Parasit., INSERM U167-CNRS 624, Institut Pasteur, 1 Rue du Pr. Calmette, 59019 Lille Cedex, France  
 SOURCE: Journal of Immunology, (1992) 148/2 (627-632).  
 ISSN: 0022-1767 CODEN: JOIMA3  
 COUNTRY: United States  
 DOCUMENT TYPE: Journal, Article  
 FILE SEGMENT: 022 Human Genetics  
 025 Hematology  
 026 Immunology, Serology and Transplantation  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB A cDNA library, constructed from purified blood eosinophils, was screened with the B cell CD23 cDNA probe. A clone designated E015 has been isolated and found to encode a non-classical HLA class I gene transcript. E015 was compared with HLA-E and found to be 99.9% similar at the nucleotide level and to extend further in the 3' untranslated region. The presence of an additional polyadenylation signal in the E015 3' end suggests that E015 clone represents a copy of the 3.3-kb mRNA species detected in Northern blot analyses. HLA-E transcripts of 1.9 and 3.3 kb have been described in a variety of cell types. The two E015 mRNA species, similar in size to the previously defined HLA-E mRNA, were present at high levels in blood leukocyte populations and at variable levels in different cell lines. The E015 transcripts were found at abundant levels in hypodense and normodense eosinophils from hypereosinophilic patients. In situ hybridization confirmed the expression of E015 mRNA in eosinophils. Neutrophils and lymphocytes from normal donors or from patients with hypereosinophilia also strongly expressed E015 mRNA. Among the cell lines studied, the highest levels of E015 transcripts were detected in B and monocytic cell lines, whereas intermediate and lower levels were found in eosinophilic, NK-like, megakaryocytic, and T cell lines, respectively. Similar to its effect on classical HLA class I transcripts, IFN-gamma increased the levels of E015 mRNA in eosinophils and neutrophils from hypereosinophilic patients. These results suggest that purified blood eosinophils as well as neutrophils express E015/HLA-E mRNA; however, further experiments are needed to investigate the localization and the function of E015 protein products.

L8 ANSWER 27 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.  
 ACCESSION NUMBER: 93075714 EMBASE  
 DOCUMENT NUMBER: 1993075714  
 TITLE: Recent advances in human natural killer cells.  
 AUTHOR: Moretta L.; Ciccone E.; Biassoni R.; Poggi A.; Mingari M.C.; Moretta A.  
 CORPORATE SOURCE: Istito Naz per la Ricerca sul Cancro, Viale Benedetto XV, 10 1-16132 Genova, Italy  
 SOURCE: International Archives of Allergy and Immunology, (1992) 99/2-4 (230-233).  
 ISSN: 1018-2438 CODEN: IAAIEG  
 COUNTRY: Switzerland  
 DOCUMENT TYPE: Journal, Conference Article  
 FILE SEGMENT: 016 Cancer  
 025 Hematology  
 026 Immunology, Serology and Transplantation  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB This paper reviews recent progresses on human Natural Killer (NK) cells which profoundly changed our concepts on NK cells and their functions. Regarding the ontogeny of NK cells, immature thymocyte populations (CD-3-4-16-) have been shown to give rise to NK cells, provided suitable culture conditions. This indicates that precursor potentially capable of differentiation towards the NK cell lineage are present within the human thymus. Although NK cells lack known receptors for antigen (i.e. sIg and TCR), NK cell populations or clones were found to be capable of mediating specific recognition of allogenic cells. This specific function was clonally distributed and, more importantly, NK clones displaying different patterns of allospecificity could be isolated from single individuals. These data indicate the existence of a NK cell repertoire for alloantigen recognition. Analysis of the surface molecules identified by NK cells indicated that certain HLA alleles (e.g. HLA-Cw3) can act as specific protective elements from lysis by clones with defined specificities (e.g. specificity 2). Therefore, HLA class I molecules appear to play a central role in the NK cell-mediated functions. The finding that human NK cells express a clonally-distributed ability to recognize alloantigens suggested the existence of distinct surface receptors. Indeed, a new family of triggering surface molecules (58 kD) has been identified by the use of monoclonal antibodies. The expression of these molecules was shown to represent a stable phenotypic property of human NK cells and to be clonally distributed. More importantly, the expression of 58 kD molecules appears to correlate with the ability to recognize given allospecificities. These data are supporting the concept that 58 kD molecules are part of receptor structures involved in the NK cell-mediated recognition.

L8 ANSWER 28 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 18  
 ACCESSION NUMBER: 90120495 EMBASE  
 DOCUMENT NUMBER: 1990120495

TITLE: Dendritic cells stimulate primary human cytolytic lymphocyte responses in the absence of CD4+ helper T cells.  
 AUTHOR: Young J.W.; Steinman R.M.  
 CORPORATE SOURCE: The Rockefeller University, Box 280, 1230 York Avenue, New York, NY 10021, United States  
 SOURCE: Journal of Experimental Medicine, (1990) 171/4 (1315-1332).  
 ISSN: 0022-1007 CODEN: JEMEA  
 COUNTRY: United States  
 DOCUMENT TYPE: Journal; Article  
 FILE SEGMENT: 025 Hematology  
 026 Immunology, Serology and Transplantation  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB Cytotoxic lymphocytes are typically generated from unfractionated suspensions of human lymphocytes by stimulating with heterogeneous APCs and exogenous growth factors. We have found that human blood dendritic cells can directly stimulate allogeneic human CD8+ T cells to proliferate and express antigen-specific cytotoxic activity. These primary responses, which are accompanied by the release of T cell growth factor(s), are induced in the absence of CD4+ helper T cells and are not inhibited by anti-CD4 mAb. Both antigen-specific CTL as well as nonspecific NK cells can be elicited by dendritic cells. The NK cell response can be depleted at the precursor level by panning with an anti-CD11b mAb, which removes a CD11b+/CD28-, CD16+ subset from the starting CD4- responders. Allogeneic blood monocytes are neither stimulatory nor inhibitory of these primary CD4- MLRs, even though monocytes present alloantigen in such a way as to be recognized as specific targets for CTL that have been sensitized by dendritic cells. The number of CD8+ cells that are blast transformed and express an activated phenotype (i.e. HLA DR/DQ+, CD25/IL-2R+, CD45R-) reaches 30-40% of the culture at day 4-5, the peak of the helper-independent response. We conclude that antigen-presentation by dendritic cells is sufficient in itself to prime cytolytic precursors. We speculate that using dendritic cell stimulators and CD4- responders in MLRs may be more efficient than standard tissue typing approaches for the detection of subtle, but important class I MHC-restricted histoincompatibilities in human transplantation.

L8 ANSWER 29 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 19  
 ACCESSION NUMBER: 89048435 EMBASE  
 DOCUMENT NUMBER: 1989048435  
 TITLE: Use of anti-HLA antibodies to mask major histocompatibility complex gene products on tumor cells can enhance susceptibility of these cells to lysis by natural killer cells.  
 AUTHOR: Lobo P.I.; Spencer C.E.  
 CORPORATE SOURCE: Department of Medicine, University of Virginia School of Medicine, Charlottesville, VA 22908, United States  
 SOURCE: Journal of Clinical Investigation, (1989) 83/1 (278-287).  
 ISSN: 0021-9738 CODEN: JCINAO  
 COUNTRY: United States  
 DOCUMENT TYPE: Journal  
 FILE SEGMENT: 016 Cancer  
 022 Human Genetics  
 026 Immunology, Serology and Transplantation  
 LANGUAGE: English  
 SUMMARY LANGUAGE: English

AB The role of major histocompatibility gene products (i.e., HLA molecules) in rendering tumor cells resistant to natural killer (NK) cell-mediated lysis was investigated by using mouse monoclonal antibodies to bind and mask HLA or non-HLA gene products on the cell membrane of human allogeneic tumor targets. Enhanced lysis of resistant lymphoid and certain other solid tumor cell lines was observed only when monoclonals used reacted to class I and II HLA molecules but not non-HLA molecules on tumor targets. Enhanced lysis was not due to antibody dependent cellular cytotoxicity or due to an effect of antibody on NK effectors. Of importance, normal autologous and allogeneic human lymphocytes could not be lysed by NK cells despite blast transformation with mitogens or masking of HLA membrane determinants on blasts with monoclonal antibodies. Enhanced lysis, in the presence of antibody to HLA antigens, was not due to increased NK cell binding to tumor targets, but a consequence of enhanced postbinding lysis. Studies using granules obtained from NK cells indicated that masking of HLA antigens did not enhance the susceptibility of tumor targets to cytotoxicity. Such observations would suggest that HLA antigens on tumor targets inhibit the triggering of effector cells (and release of cytotoxicins) after recognition and binding of NK cells to target cells.

L8 ANSWER 30 OF 30 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 20  
 ACCESSION NUMBER: 88000518 EMBASE  
 DOCUMENT NUMBER: 1988000518  
 TITLE: [Bronchioloalveolar lavage in conjunction with transbronchial lung biopsy: Value and indications]. AUSSAGE- UND EINSAATZMOGLICHKEITEN DER BRONCHIOLOALVEOLAREN LAVAGE KOMBINIERT MIT TRANSBRONCHIALER LUNGENBIOPSIE.  
 AUTHOR: Popper H.; Pongratz M.  
 CORPORATE SOURCE: Laboratorium für Umwelt- und Atemtraktpathologie, Institut für Pathologische Anatomie, Universität Graz, 8036 Graz, Austria  
 SOURCE: Wiener Klinische Wochenschrift, (1987) 99/24 (848-855).  
 ISSN: 0043-5325 CODEN: WKWOAO  
 COUNTRY: Austria  
 DOCUMENT TYPE: Journal  
 FILE SEGMENT: 005 General Pathology and Pathological Anatomy  
 015 Chest Diseases, Thoracic Surgery and Tuberculosis  
 LANGUAGE: German  
 SUMMARY LANGUAGE: English

AB Bronchioloalveolar lavage (BAL) enables diffuse interstitial lung disease to be divided into lymphocytic and granulocytic alveolitis. The combination of BAL and transbronchial lung biopsies using modern flexible fiberoptic bronchoscopes allows the subdivision of lymphocytic alveolitis into sarcoidosis, exogenous allergic alveolitis (synonym: hypersensitivity pneumonitis; EAA) and granulomatous pneumonias caused by infectious agents. The use of immunohistochemical surface markers of lymphocytes in conjunction with BAL provides further differentiation of lymphocytes into T- and B-, T-helper and T-suppressor types, natural killer cells (NK cells) and cytotoxic T-cells. A predominance of T-suppressor lymphocytes is an indication of EAA, whereas a predominance of T-helper

lymphocytes is positively correlated with sarcoidosis. Other markers, e.g. HLA-DR, when expressed on the surface of alveolar macrophages, merely indicate activation unrelated to a specific type of lymphocytic alveolitis. BAL is also a new and promising diagnostic tool for pneumoconioses and other types of lung disease caused by inhaled industrial pollutants. Ferruginous bodies and silica crystals, free or ingested by alveolar macrophages, can be found more easily than by scraping tissue blocks or from multiple sections of transbronchial biopsies. BAL cells can easily be processed for electron microscopy and inhaled foreign material can be analysed in an electron microscope using X-ray diffraction analysis (EDAX) or electron spectroscopic imaging (ESI). BAL is also of value in the diagnosis of peripheral lung carcinomas, in addition to cytologic sputum analysis, brush smears, transthoracic fine needle aspiration and transbronchial biopsies. BAL is a valuable diagnostic tool in cases of unusual pneumonia where fungi can be visualized by silver impregnation techniques and viruses by antibodies using immunofluorescence microscopy.

=> dis his

(FILE 'HOME' ENTERED AT 13:53:22 ON 24 JUN 2002)

FILE 'MEDLINE, CAPLUS, EMBASE, BIOSIS' ENTERED AT 13:53:34 ON 24 JUN 2002

```

L1 3511 S BRAUD V?/AU OR ALLAN D?/AU OR OGG G?/AU OR OCALLAGHAN C?/AU O
L2 47 S L1 AND (NK?)
L3 35 S L2 AND (HLA (1N) E)
L4 14 DUP REM L3 (21 DUPLICATES REMOVED)
L5 72958 S NK (P) (NK? OR CD94?)
L6 315 S (HLA (1N) E) (P) (NK? OR CD94?)
L7 63 S L6 AND PD<19981204
L8 30 DUP REM L7 (33 DUPLICATES REMOVED)

```

(FILE 'HOME' ENTERED AT 13:53:22 ON 24 JUN 2002)

FILE 'MEDLINE, CAPLUS, EMBASE, BIOSIS' ENTERED AT 13:53:34 ON 24 JUN 2002

L1 3511 S BRAUD V?/AU OR ALLAN D?/AU OR OGG G?/AU OR OCALLAGHAN C?/AU O  
L2 47 S L1 AND (NK?)  
L3 35 S L2 AND (HLA (1N) E)  
L4 14 DUP REM L3 (21 DUPLICATES REMOVED)  
L5 72958 S NK (P) (NK? OR CD94?)  
L6 315 S (HLA (1N) E) (P) (NK? OR CD94?)  
L7 63 S L6 AND PD<19981204  
L8 30 DUP REM L7 (33 DUPLICATES REMOVED)